

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



INTERNATIONAL BUREAU OF PATENT COOPERATION  
358, RUE DE LA PATENTE  
CH-1202 GENEVE 20  
SUISSE

(43) International Publication Date  
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number  
**WO 02/053594 A2**

- (51) International Patent Classification<sup>7</sup>: C07K 14/81, A61P 25/28, A61K 38/55, C07K 5/03
- (21) International Application Number: PCT/US01/50826
- (22) International Filing Date:  
28 December 2001 (28.12.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/258,705 28 December 2000 (28.12.2000) US  
60/275,756 14 March 2001 (14.03.2001) US
- (71) Applicants (for all designated States except US): OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 N.E. 13th Street, Oklahoma City, OK 73104 (US). THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ILLINOIS [US/US]; 201 South Wright Street, 352 Henry Administration Building, Urbana, IL 60101 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): TANG, Jordan, J., N. [US/US]; 1204 Leawood Drive, Edmond, OK 73034 (US). KOELSCH, Gerald [US/US]; 8317 N.W. 105 Terrace, Oklahoma City, OK 73162 (US). GHOSH, Arun, K. [US/US]; 1407 Clinton Place, River Forest, IL 60305 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**BEST AVAILABLE COPY**

(54) Title: INHIBITORS OF MEMAP SIN 2 AND USE THEREOF

(57) Abstract: Methods for the production of purified, catalytically active, recombinant memapsin 2 have been developed. The substrate and subsite specificity of the catalytically active enzyme have been determined by a method which determines the initial hydrolysis rate of the substrate by using MALDI-TOF/MS. Alternatively, the subsite specificity of memapsin can be determined by probing a library of inhibitors with memapsin 2 and subsequently detecting the bound memapsin 2 with an antibody raised to memapsin 2 and an alkaline phosphatase conjugated secondary antibody. The substrate and subsite specificity information was used to design substrate analogs of the natural memapsin 2 substrate that can inhibit the function of memapsin2. The substrate analogs are based on peptide sequences, shown to be related to the natural peptide substrates for memapsin 2. The substrate analogs contain at least one analog of an amide bond which is not capable of being cleaved by memapsin 2. Processes for the synthesis of substrate analogues including isosteres at the sites of the critical amino acid residues were developed and the more than seventy substrate analogues were synthesized, among which MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-066, MMI-070, and MMI-071 have inhibition constants in the range of  $1.4-61.4 \times 10^9$  M against recombinant pro-memapsin 2. These inhibitors are useful in diagnostics and for the treatment and/or prevention of Alzheimer's disease.

WO 02/053594 A2

B/3

## INHIBITORS OF MEMAP SIN 2 AND USE THEREOF

## RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Nos. 60/275,756, filed on March 14, 2001 and 60/258,705 filed on December 28, 2000,  
5 the entire teachings of both of which are incorporated by reference in their entirety.

## BACKGROUND OF THE INVENTION

This invention is in the area of the design and synthesis of specific inhibitors of the aspartic protease Memapsin 2 (beta-secretase, or  $\beta$ -secretase) which are useful in the treatment and/or prevention of Alzheimer's Disease.

10 Alzheimer's disease (AD) is a degenerative disorder of the brain first described by Alios Alzheimer in 1907 after examining one of his patients who suffered drastic reduction in cognitive abilities and had generalized dementia (*The early story of Alzheimer's Disease*, edited by Bick *et al.* (Raven Press, New York 1987)). It is the leading cause of dementia in elderly persons. AD patients have  
15 increased problems with memory loss and intellectual functions which progress to the point where they cannot function as normal individuals. With the loss of intellectual skills the patients exhibit personality changes, socially inappropriate actions and schizophrenia (*A Guide to the Understanding of Alzheimer's Disease and Related Disorders*, edited by Jorm (New York University Press, New York  
20 1987)). AD is devastating for both victims and their families, for there is no effective palliative or preventive treatment for the inevitable neurodegeneration.

AD is associated with neuritic plaques measuring up to 200  $\mu$ m in diameter in the cortex, hippocampus, subiculum, hippocampal gyrus, and amygdala. One of the principal constituents of neuritic plaques is amyloid, which is stained by Congo  
25 Red (Fisher (1983); Kelly Microbiol. Sci. 1(9):214-219 (1984)). Amyloid plaques stained by Congo Red are extracellular, pink or rust-colored in bright field, and birefringent in polarized light. The plaques are composed of polypeptide fibrils and

are often present around blood vessels, reducing blood supply to various neurons in the brain.

Various factors such as genetic predisposition, infectious agents, toxins, metals, and head trauma have all been suggested as possible mechanisms of AD neuropathy. Available evidence strongly indicates that there are distinct types of genetic predispositions for AD. First, molecular analysis has provided evidence for mutations in the amyloid precursor protein (APP) gene in certain AD-stricken families (Goate *et al.* *Nature* 349:704-706 (1991); Murrell *et al.* *Science* 254:97-99 (1991); Chartier-Harlin *et al.* *Nature* 353:844-846 (1991); Mullan *et al.*, *Nature Genet.* 1:345-347 (1992)). Additional genes for dominant forms of early onset AD reside on chromosome 14 and chromosome 1 (Rogaev *et al.*, *Nature* 376:775-778 (1995); Levy-Lahad *et al.*, *Science* 269:973-977 (1995); Sherrington *et al.*, *Nature* 375:754-760 (1995)). Another loci associated with AD resides on chromosome 19 and encodes a variant form of apolipoprotein E (Corder, *Science* 261:921-923 (1993)).

Amyloid plaques are abundantly present in AD patients and in Down's Syndrome individuals surviving to the age of 40. The overexpression of APP in Down's Syndrome is recognized as a possible cause of the development of AD in Down's patients over thirty years of age (Rumble *et al.*, *New England J. Med.* 320:1446-1452 (1989); Mann *et al.*, *Neurobiol. Aging* 10:397-399 (1989)). The plaques are also present in the normal aging brain, although at a lower number. These plaques are made up primarily of the amyloid  $\beta$  peptide ( $A\beta$ ; sometimes also referred to in the literature as  $\beta$ -amyloid peptide or  $\beta$  peptide) (Glenner and Wong, *Biochem. Biophys. Res. Comm.* 120:885-890 (1984)), which is also the primary protein constituent in cerebrovascular amyloid deposits. The amyloid is a filamentous material that is arranged in beta-pleated sheets.  $A\beta$  is a hydrophobic peptide comprising up to 43 amino acids.

The determination of its amino acid sequence led to the cloning of the APP cDNA (Kang *et al.*, *Nature* 325:733-735 (1987); Goldgaber *et al.*, *Science* 235:877-880 (1987); Robakis *et al.*, *Proc. Natl. Acad. Sci.* 84:4190-4194 (1987); Tanzi *et al.*, *Nature* 331:528-530 (1988)) and genomic APP DNA (Lemaire *et al.*, *Nucl. Acids*

Res. 17:517-522 (1989); Yoshikai *et al.*, *Gene* 87, 257-263 (1990)). A number of forms of APP cDNA have been identified, including the three most abundant forms, APP695, APP751, and APP770. These forms arise from a single precursor RNA by alternate splicing. The gene spans more than 175 kb with 18 exons (Yoshikai *et al.* 5 (1990)). APP contains an extracellular domain, a transmembrane region and a cytoplasmic domain. A $\beta$  consists of up to 28 amino acids just outside the hydrophobic transmembrane domain and up to 15 residues of this transmembrane domain. A $\beta$  is normally found in brain and other tissues such as heart, kidney and spleen. However, A $\beta$  deposits are usually found in abundance only in the brain.

10 Van Broeckhoven *et al.*, *Science* 248:1120-1122 (1990), have demonstrated that the APP gene is tightly linked to hereditary cerebral hemorrhage with amyloidosis (HCHWA-D) in two Dutch families. This was confirmed by the finding of a point mutation in the APP coding region in two Dutch patients (Levy *et al.*, *Science* 248:1124-1128 (1990)). The mutation substituted a glutamine for glutamic acid at position 22 of the A $\beta$  (position 618 of APP695, or position 693 of APP770). 15 In addition, certain families are genetically predisposed to Alzheimer's disease, a condition referred to as familial Alzheimer's disease (FAD), through mutations resulting in an amino acid replacement at position 717 of the full length protein (Goate *et al.* (1991); Murrell *et al.* (1991); Chartier-Harlin *et al.* (1991)). These 20 mutations co-segregate with the disease within the families and are absent in families with late-onset AD. This mutation at amino acid 717 increases the production of the A $\beta$ <sub>1-42</sub> form of A $\beta$  from APP (Suzuki *et al.*, *Science* 264:1336-1340 (1994)). Another mutant form contains a change in amino acids at positions 670 and 671 of the full length protein (Mullan *et al.* (1992)). This mutation to 25 amino acids 670 and 671 increases the production of total A $\beta$  from APP (Citron *et al.*, *Nature* 360:622-674 (1992)).

APP is processed *in vivo* at three sites. The evidence suggests that cleavage at the  $\beta$ -secretase site by a membrane associated metalloprotease is a physiological event. This site is located in APP 12 residues away from the luminal surface of the 30 plasma membrane. Cleavage of the  $\beta$ -secretase site (28 residues from the plasma membrane's luminal surface) and the  $\beta$ -secretase site (in the transmembrane region)

results in the 40/42-residue  $\beta$ -amyloid peptide ( $A\beta$ ), whose elevated production and accumulation in the brain are the central events in the pathogenesis of Alzheimer's disease (for review, see Selkoe, D.J. *Nature* 399:23-31 (1999)). Presenilin 1, another membrane protein found in human brain, controls the hydrolysis at the APP  $\gamma$   $\beta$ -secretase site and has been postulated to be itself the responsible protease (Wolfe, M.S. *et al.*, *Nature* 398:513-517 (1999)). Presenilin 1 is expressed as a single chain molecule and its processing by a protease, presenilinase, is required to prevent it from rapid degradation (Thinakaran, G. *et al.*, *Neuron* 17:181-190 (1996) and Podlisny, M.B., *et al.*, *Neurobiol. Dis.* 3:325-37 (1997)). The identity of presenilinase is unknown. The *in vivo* processing of the  $\beta$ -secretase site is thought to be the rate-limiting step in  $A\beta$  production (Sinha, S. & Lieberburg, I., *Proc. Natl. Acad. Sci., USA*, 96:11049-11053 (1999)), and is therefore a strong therapeutic target.

The design of inhibitors effective in decreasing amyloid plaque formation is dependent on the identification of the critical enzyme(s) in the cleavage of APP to yield the 42 amino acid peptide, the  $A\beta_{1-42}$  form of  $A\beta$ . Although several enzymes have been identified, it has not been possible to produce active enzyme. Without active enzyme, one cannot confirm the substrate specificity, determine the subsite specificity, nor determine the kinetics or critical active site residues, all of which are essential for the design of inhibitors.

Memapsin 2 has been shown to be beta-secretase, a key protease involved in the production in human brain of beta-amyloid peptide from beta-amyloid precursor protein (for review, see Selkoe, D.J. *Nature* 399:23-31 (1999)). It is now generally accepted that the accumulation of beta-amyloid peptide in human brain is a major cause for Alzheimer's disease. Inhibitors specifically designed for human memapsin 2 should inhibit or decrease the formation of beta-amyloid peptide and the progression of the Alzheimer's disease.

Memapsin 2 belongs to the aspartic protease family. It is homologous in amino acid sequence to other eukaryotic aspartic proteases and contains motifs specific to that family. These structural similarities predict that memapsin 2 and other eukaryotic aspartic proteases share common catalytic mechanism, Davies,

-5-

D.R., *Annu. Rev. Biophys. Chem.* 19, 189 (1990). The most successful inhibitors for aspartic proteases are mimics of the transition state of these enzymes. These inhibitors have substrate-like structure with the cleaved planar peptide bond between the carbonyl carbon and the amide nitrogen replaced by two tetrahedral  
5 atoms, such as hydroxyethylene [-CH(OH)-CH<sub>2</sub>-], which was originally discovered in the structure of pepstatin (Marciniszyn *et al.*, 1976).

A need exists to develop new, improved inhibitors of proteases involved in the production of beta-amyloid protein from beta-amyloid precursor protein, such as memapsin 2 inhibitors, that are effective, for example, in the treatment of  
10 Alzheimer's disease in humans.

#### SUMMARY OF THE INVENTION

The present invention relates to inhibitors of memapsin 2 activity and methods of using the inhibitors of memapsin 2 to treat Alzheimer's disease in humans.

15 In one embodiment, the invention is an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an K<sub>i</sub> of less than or equal to 10<sup>-7</sup> M.

In another embodiment, the invention includes a compound selected from the  
20 group consisting of MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-066, MMI-070, and MMI-071.

In yet another embodiment, the invention includes a compound selected from the group consisting of MMI-012, MMI-017, MMI-018, MMI-026, MMI-037, MMI-  
25 039, MMI-040, MMI-070 and MMI-071.

Another embodiment includes a method for treating a patient to decrease the likelihood of developing or the progression of Alzheimer's disease comprising administering to the individual an effective amount of an inhibitor of memapsin 2 selected from the group consisting of MMI-005, MMI-012, MMI-017, MMI-018,

-6-

MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-070 and MMI-071.

In an additional embodiment, the invention is a method of determining the substrate side-chain preference in memapsin 2 sub-sites, comprising the steps of  
5 reacting a mixture of memapsin 2 substrates with memapsin 2 and determining the sub-site preference of memapsin 2 by determining relative initial hydrolysis rates of the mixture of memapsin 2 substrates.

In still another embodiment, the invention includes a method of determining the substrate side-chain preference in memapsin 2 sub-sites. A combinatorial library  
10 of memapsin 2 inhibitors wherein the inhibitors comprise a base sequence taken from OM99-2 is prepared. The library of inhibitors is probed with memapsin 2 wherein the memapsin 2 may bind one or a plurality of inhibitors to generate one or a plurality of bound memapsin 2 and the bound memapsin 2 is detected with an antibody raised to memapsin 2 and an alkaline phosphatase conjugated secondary  
15 antibody.

In a further embodiment, the invention includes a method of treating a human suffering from Alzheimer's disease comprising administering to the human an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate  
20 binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M.

In yet another embodiment, the invention relates to a method of treating a human suffering from Alzheimer's disease comprising administering to the human an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate  
25 binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M, wherein the inhibitor has a root mean square difference of less than or equal to 0.5 Å for the side chain and backbone atoms for amino acids 28-441 of SEQ ID NO: 2.

An additional embodiment of the invention is a method of treating a human suffering from Alzheimer's disease comprising administering to the human a  
30 compound selected from the group consisting of MMI-012, MMI-017, MMI-018, MMI-026, MMI-037, MMI-039, MMI-040, MMI-070 and MMI-071.

-7-

In yet another embodiment, the invention relates to the use of an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M for the manufacture of a  
5 medicament for the treatment of Alzheimer's disease in a human.

In an additional embodiment, the invention relates to the use of an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M, wherein the inhibitor has a  
10 root mean square difference of less than or equal to 0.5 Å for the side chain and backbone atoms for amino acids 18-379 of memapsin 2, for the manufacture of a medicament for the treatment of Alzheimer's disease in a human.

In still another embodiment, the invention relates to the use of a compound selected from the group consisting of MMI-005, MMI-012, MMI-017, MMI-018,  
15 MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-066, MMI-070 and MMI-071, for the manufacture of a medicament for the treatment of Alzheimer's disease in a human.

The substrate and subsite specificity of recombinant, catalytically active memapsin 2 was used to design substrate analogs of the natural memapsin 2  
20 substrate that can inhibit the function of memapsin 2. Initially, X-ray crystallography of memapsin 2 bound to a substrate analog, OM99-2, was used to determine the three dimensional structure of the memapsin 2, as well as the importance of the various residues in binding. Substrate analogs were then designed based on peptide sequences shown to be related to the natural peptide substrates for  
25 memapsin 2 and the crystallographic structure. The substrate analogs contain at least one analog of an amide (peptide) bond that is not capable of being cleaved by memapsin 2.

The substrate and subsite specificity of the catalytically active memapsin 2 have been further determined by a method which determines the initial hydrolysis  
30 rate of the substrates using mass spectroscopy, for example, MALDI-TOF/MS (matrix assisted laser desorption/ionization-time of flight /mass spectroscopy).



-8-

Alternatively, the subsite specificity of memapsin was further determined by probing a library of inhibitors with memapsin 2 and subsequently detecting the bound memapsin 2 with an antibody raised to memapsin 2 and an alkaline phosphatase conjugated secondary antibody. The substrate and subsite specificity information  
5 was used to design additional substrate analogs of the natural memapsin 2 substrate that can inhibit the function of memapsin 2.

Processes for the synthesis of substrate analogues including isosteres at the sites of the critical amino acid residues were developed. Substrate analogues, OM99-1, OM99-2 and more than seventy other substrate analogues were  
10 synthesized. OM99-2 is based on an octapeptide Glu-Val-Asn-Leu-Ala-Ala-Glu-Phe (SEQ ID NO: 28) with the Leu-Ala peptide bond substituted by a transition-state isostere hydroxyethylene group. The inhibition constant of OM99-2 is  $1.6 \times 10^{-9}$  M against recombinant pro-memapsin 2. The inhibition constants of MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040,  
15 MMI-066, MMI-070, and MMI-071 have inhibition constants in the range of  $1.4 - 61.4 \times 10^{-9}$  M against recombinant pro-memapsin 2.

Compositions that inhibit memapsin 2 aspartic protease activity can be small molecules, which readily pass across the blood brain barrier, are administered orally, and are not inactivated by intestinal enzymes. Furthermore, it is desirable that such  
20 compositions are relatively inexpensive to manufacture and preferentially inhibit memapsin 2 cleavage of beta-amyloid precursor protein.

This information can be used by those skilled in the art to design additional new inhibitors, using commercially available software programs and techniques familiar to those in organic chemistry and enzymology, to design new inhibitors.  
25 For example, the side chains of the inhibitors may be modified to produce stronger interactions (through hydrogen bonding, hydrophobic interaction, charge interaction and/or van der Waal interaction) in order to increase inhibition potency. Based on this type of information, the residues with minor interactions may be eliminated from the new inhibitor design to decrease the molecular weight of the inhibitor. The  
30 side chains with no structural hindrance from the enzyme may be cross-linked to lock in the effective inhibitor conformation. This type of structure also enables the

-9-

design of peptide surrogates which may effectively fill the binding sites of memapsin 2 yet produce better pharmaceutical properties. The rational design and screening of compounds for inhibitors, and their characterization are provided in this invention. Compositions effective for inhibition of memapsin 2 include small molecule inhibitors, and inhibitors that are capable of crossing the blood brain barrier. Such inhibitors can interact with memapsin 2, or its substrate, to inhibit cleavage by memapsin 2.

The invention described herein provides compounds which inhibit memapsin 2 activity, in particular the aspartic protease activity of memapsin 2, which converts beta-amyloid precursor protein to beta-amyloid protein. The compounds of the invention can be used to treat Alzheimer's disease in humans.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of inhibitors for memapsin 2. "A" and "B" denote aliphatic linkages (saturated or partially unsaturated) of any number of carbons, between side chains in positions  $P_1$  and  $P_3$ , for example, the amino acid side chains  $P_1$  Leu and  $P_3$  Val in these respective positions. Other structural elements of the inhibitor are omitted for clarity.

Figure 2 is a schematic representation of inhibitors for memapsin 2. "A" denotes aliphatic linkages (saturated or partially unsaturated) of any number of carbons, between side chains in positions  $P_2$  and  $P_4$ , for example, the amino acid side chains  $P_2$  Asn and  $P_4$  Glu in these respective positions. Other structural elements of the inhibitor are omitted for clarity.

Figure 3 is a schematic of the design for the side chain at the  $P_1'$  subsite for the new memapsin 2 inhibitors based on the current crystal structure. Arrows indicate possible interactions between memapsin 2 and inhibitor. Other structural elements of inhibitor are omitted for clarity.

Figure 4 is a schematic representation of inhibitors for memapsin 2. "A" and "B" denote aliphatic linkages (saturated or partially unsaturated) between side chains in positions  $P_1$  and the backbone atoms of  $P_3$ , for example, the amino acid

-10-

side chains P<sub>1</sub> Leu in this position. Other structural elements of the inhibitor are omitted for clarity.

Figure 5 shows the relative specificity of memapsin 2 for amino acid residues in positions P<sub>1</sub>'-P<sub>4</sub>'. Letters above bars indicate the native amino acid at that position. Catalytic efficiency is expressed relative to the native amino acid at each position.

Figure 6 depicts the nucleotide sequence of human Memapsin 2 (SEQ ID NO: 1).

Figures 7A and 7B depict the partial protein sequence of human Memapsin 2, excluding the signal peptide (SEQ ID NO: 2). Amino acids 28-48 are remnant putative propeptide residues. Amino acids 58-61, 78, 80, 82-83, 116, 118-121, 156, 166, 174, 246, 274, 276, 278-281, 283, and 376-377 are residues in contact with the OM99-2 inhibitor. Amino acids 54-57, 61-68, 73-80, 86-89, 109-111, 113-118, 123-134, 143-154, 165-168, 198-202, and 220-224 are N-lobe beta strands. Amino acids 184-191 and 210-217 are N-lobe helices. Amino acids 237-240, 247-249, 251-256, 259-260, 273-275, 282-285, 316-318, 331-336, 342-348, 354-357, 366-370, 372-375, 380-383, 390-395, 400-405, and 418-420 are C-lobe beta strands. Amino acids 286-299, 307-310, 350-353, 384-387, and 427-431 are C-lobe helices.

Figures 8A and 8B depict the protein sequence of human promemapsin 2 (SEQ ID NO: 3). Amino acids 1-15 are vector-derived residues. Amino acids 16-63 are a putative pro peptide. Amino acids 1-13 are derived from the T7 promoter. Amino acids 16-456 are Pro-memapsin 2-T1. Amino acids 16-421 are Promemapsin 2-T2.

Figure 9 depicts the amino acid sequence of human pre-promemapsin 2 (SEQ ID NO: 4). Amino acids 1-13 are the signal peptide. Amino acids 41-454 correspond to amino acids 28-441 of Figures 7A and 7B, and amino acids 43-456 of Figures 8A and 8B. The active site aspartic acids are at amino acid positions 93 and 289.

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention.

-11-

## DETAILED DESCRIPTION OF THE INVENTION

The features and other details of the invention, either as steps of the invention or as combinations of parts of the invention, will now be more particularly described and pointed out in the claims. It will be understood that the particular  
 5 embodiments of the invention are shown by way of illustration and not as limitations of the invention. The principle features of this invention can be employed in various embodiments without departing from the scope of the invention.

**I. Design and Synthesis of Inhibitors****Design of Substrate Analogs for Memapsin 2.**

10 The five human aspartic proteases have homologous amino acid sequences and have similar three-dimensional structures. There are two aspartic residues in the active site and each residue is found within the signature aspartic protease sequence motif, Asp-Thr/Ser-Gly-. There are generally two homologous domains within an aspartic protease and the substrate binding site is positioned between these two  
 15 domains, based on the three-dimensional structures. The substrate binding sites of aspartic proteases generally recognize eight amino acid residues. There are generally four residues on each side of the amide bond that is cleaved by the aspartic protease.

Typically the side chains of each amino acid are involved in the specificity of the substrate/aspartic protease interaction. The side chain of each substrate residue  
 20 is recognized by regions of the enzyme which are collectively called sub-sites. The generally accepted nomenclature for the protease sub-sites and their corresponding substrate residues are shown below, where the double slash represents the position of bond cleavage.

Protease sub-sites	S4	S3	S2	S1	S1'	S2'	S3'	S4'
25 Substrate residues	P4	P3	P2	P1	// P1'	P2'	P3'	P4'

While there is a general motif for aspartic protease substrate recognition, each protease has a very different substrate specificity and breadth of specificity.

-12-

Once the specificity of an aspartic protease is known, inhibitors can be designed based on that specificity, which interact with the aspartic protease in a way that prevents natural substrate from being efficiently cleaved. Some aspartic proteases have specificities that can accommodate many different residues in each of the sub-  
5 sites for successful hydrolysis. Pepsin and cathepsin D have this type of specificity and are said to have "broad" substrate specificity. When only a very few residues can be recognized at a sub-site, such as in renin, the aspartic protease is said to have a stringent or narrow specificity.

The information on the specificity of an aspartic protease can be used to  
10 design specific inhibitors in which the preferred residues are placed at specific sub-sites and the cleaved peptide bond is replaced by an analog of the transition-state. These analogs are called transition state isosteres. Aspartic proteases cleave amide bonds by a hydrolytic mechanism. This reaction mechanism involves the attack by a hydroxide ion on the  $\beta$ -carbon of the amino acid. Protonation must occur at the  
15 other atom attached to the  $\beta$ -carbon through the bond that is to be cleaved. If the  $\beta$ -carbon is insufficiently electrophilic or the atom attached to the bond to be cleaved is insufficiently nucleophilic, the bond will not be cleaved by a hydrolytic mechanism. Analogs exist which do not mimic the transition state but which are non-hydrolyzable, but transition state isosteres mimic the transition state specifically and  
20 are non-hydrolyzable.

Transition state theory indicates that it is the transition state intermediate of the reaction which the enzyme catalyzes for which the enzyme has its highest affinity. It is the transition state structure, not the ground state structure, of the substrate which will have the highest affinity for its given enzyme. The transition  
25 state for the hydrolysis of an amide bond is tetrahedral while the ground state structure is planar. A typical transition-state isostere of aspartic protease is -CH(OH)-CH<sub>2</sub>-, as was first discovered in pepstatin by Marciniszyn *et al.* (1976). The transition-state analogue principles have been successfully applied to inhibitor drugs for human immunodeficiency virus protease, an aspartic protease. Many of  
30 these are currently in clinical use. Information on the structure, specificity, and types of inhibitors can be found in Tang, Acid Proteases, Structure, Function and Biology,

-13-

Adv. in Exptl. Med. Biol. vol. 95 (Plenum Press, NY 1977); Kostka, Aspartic Proteinases and their Inhibitors (Walter de Gruyter, Berlin 1985); Dunn, Structure and Functions of the Aspartic Proteinases, Adv. in Exptl. Med. Biol. 306 (Plenum Press, NY 1991); Takahashi, Aspartic Proteases, Structure, Function, Biology, Biomedical Implications, Adv. in Exptl. Med. Biol. 362 (Plenum Press, NY 1995); and James, Aspartic Proteinases, Retroviral and Cellular Enzymes, Adv. in Exptl. Med. Biol. 436 (Plenum Press, NY 1998), the teachings of all of which are incorporated herein in their entirety).

Substrate analog compositions are generally of the general formula  $X-L_4-P_4-L_3-P_3-L_2-P_2-L_1-P_1-L_0-P_1'-L_1'-P_2'-L_2'-P_3'-L_3'-P_4'-L_4'-Y$ . The substrate analog compositions are analogs of small peptide molecules. Their basic structure is derived from peptide sequences that were determined through structure/function studies. It is understood that positions represented by  $P_x$  represent the substrate specificity position relative to the cleavage site which is represented by an  $-L_0-$ . The positions of the compositions represented by  $L_x$  represent the linking regions between each substrate specificity position,  $P_x$ .

In a natural substrate for memapsin 2, a  $P_x-L_x$  pair would represent a single amino acid of the peptide which is to be cleaved. In the present general formula, each  $P_x$  part of the formula refers to the  $\alpha$ -carbon and side chain functional group of each would be amino acid. Thus, the  $P_x$  portion of an  $P_x-L_x$  pair for alanine represents  $HC-CH_3$ . The general formula representing the  $P_x$  portion of the general composition is  $-R_1CR_3-$ .

In general  $R_1$  can be either  $CH_3$  (side chain of alanine),  $CH(CH_3)_2$  (side chain of valine),  $CH_2CH(CH_3)_2$  (side chain of leucine),  $(CH_3)CH(CH_2CH_3)$  (side chain of isoleucine),  $CH_2$ (indole) (side chain of tryptophan),  $CH_2$ (benzene) (side chain of phenylalanine),  $CH_2CH_2SCH_3$  (side chain of methionine),  $H$  (side chain of glycine),  $CH_2OH$  (side chain of serine),  $CHOHCH_3$  (side chain of threonine),  $CH_2$ (phenol) (side chain of tyrosine),  $CH_2SH$  (side chain of cysteine),  $CH_2CH_2CONH_2$  (side chain of glutamine),  $CH_2CONH_2$  (side chain of asparagine),  $CH_2CH_2CH_2CH_2NH_2$  (side chain of lysine),  $CH_2CH_2CH_2NHC(NH)(NH_2)$  (side chain of arginine),  $CH_2$ (imidazole) (side chain of histidine),  $CH_2COOH$  (side chain of aspartic acid),

-14-

$\text{CH}_2\text{CH}_2\text{COOH}$  (side chain of glutamic acid), and functional natural and non-natural derivatives or synthetic substitutions of these.

It is most preferred that  $\text{R}_3$  is a single H. In general, however,  $\text{R}_3$  can be alkenyl, alkynyl, alkenyloxy, and alkynyloxy groups that allow binding to memapsin

5 2. Preferably, alkenyl, alkynyl, alkenyloxy and alkynyloxy groups have from 2 to 40 carbons, and more preferably from 2 to 20 carbons, from 2 to 10 carbons, or from 2 to 3 carbons, and functional natural and non-natural derivatives or synthetic substitutions of these.

The  $\text{L}_x$  portion of the  $\text{P}_x\text{-L}_x$  pair represents the atoms linking the  $\text{P}_x$  regions  
10 together. In a natural substrate the  $\text{L}_x$  represents the  $\beta$ -carbon attached to the amino portion of what would be the next amino acid in the chain. Thus,  $\text{L}_x$  would be represented by  $-\text{CO-NH}-$ . The general formula for  $\text{L}_x$  is represented by  $\text{R}_2$ . In general  $\text{R}_2$  can be  $\text{CO-NH}$  (amide),  $\text{CH}(\text{OH})(\text{CH}_2)$  (hydroxyethylene),  $\text{CH}(\text{OH})\text{CH}(\text{OH})$  (dihydroxyethylene),  $\text{CH}(\text{OH})\text{CH}_2\text{NH}$  (hydroxyethylamine),  
15  $\text{PO}(\text{OH})\text{CH}_2$  (phosphinate),  $\text{CH}_2\text{NH}$  (reduced amide). It is understood that more than one  $\text{L}_x$  maybe an isostere as long as the substrate analog functions to inhibit aspartic protease function.

$\text{L}_x$ s which are not isosteres may either be an amide bond or mimetic of an amide bond that is non-hydrolyzable.

20 X and Y represent molecules which are not typically involved in the recognition by the aspartic protease recognition site, but which do not interfere with recognition. It is preferred that these molecules confer resistance to the degradation of the substrate analog. Preferred examples would be amino acids coupled to the substrate analog through a non-hydrolyzable bond. Other preferred compounds are  
25 capping agents. Still other preferred compounds are compounds that could be used in the purification of the substrate analogs such as biotin.

As used herein, alkyl refers to substituted or unsubstituted straight, branched or cyclic alkyl groups; and alkoxy refers to substituted or unsubstituted straight, branched or cyclic alkoxy. Preferably, alkyl and alkoxy groups have from 1  
30 to 40 carbons, and more preferably from 1 to 20 carbons, from 1 to 10 carbons, or from 1 to 3 carbons.

-15-

As used herein, alkenyl refers to substituted or unsubstituted straight chain or branched alkenyl groups; alkynyl refers to substituted or unsubstituted straight chain or branched alkynyl groups; alkenyloxy refers to substituted or unsubstituted straight chain or branched alkenyloxy; and alkynyloxy refers to substituted or unsubstituted straight chain or branched alkynyloxy. Preferably, alkenyl, alkynyl, alkenyloxy and alkynyloxy groups have from 2 to 40 carbons, and more preferably from 2 to 20 carbons, from 2 to 10 carbons, or from 2 to 3 carbons.

As used herein, alkaryl refers to an alkyl group that has an aryl substituent; aralkyl refers to an aryl group that has an alkyl substituent; heterocyclic-alkyl refers to a heterocyclic group with an alkyl substituent; alkyl-heterocyclic refers to an alkyl group that has a heterocyclic substituent.

The substituents for alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, and alkynyloxy groups can be halogen, cyano, amino, thio, carboxy, ester, ether, thioether, carboxamide, hydroxy, or mercapto. Further, the groups can optionally have one or more methylene groups replaced with a heteroatom, such as O, NH or S.

A number of different substrates were tested and analyzed, and the cleavage rules for Memapsin 2 were determined. The results of the substrates which were analyzed are presented in Table 1 and the rules determined from these results are summarized below.

(1) The primary specificity site for a memapsin 2 substrate is subsite position,  $S_1'$ . This means that the most important determinant for substrate specificity in memapsin 2 is the amino acid,  $P_1'$ .  $P_1'$  may be a small side chain for memapsin 2 to recognize the substrate. Preferred embodiments are substrate analogs where  $R_1$  of the  $P_1'$  position is either H (side chain of glycine),  $CH_3$  (side chain of alanine),  $CH_2OH$  (side chain of serine), or  $CH_2COOH$  (side chain of aspartic acid). Embodiments that have an  $R_1$  structurally smaller than  $CH_3$  (side chain of alanine) or  $CH_2OH$  (side chain of serine) are also preferred. However, substrates in Table 1 may not, due to their amino acid composition, provide a complete representation of the substrate specificity of memapsin 2. Therefore,  $P_1'$  may not be limited to the small residues mentioned, but may also include the following:



-16-

- $\text{CH}_2\text{CH}(\text{CH}_3)_2$  (sidechain of leucine),  
 $(\text{CH}_3)\text{CH}(\text{CH}_2\text{CH}_3)$  (sidechain of isoleucine),  
 $\text{CH}_2(\text{INDOLE})$  (sidechain of tryptophan),  
 $\text{CH}_2(\text{BENZENE})$  (sidechain of phenylalanine),  
5  $\text{CH}(\text{CH}_3)_2$  (sidechain of valine),  
 $\text{CH}_2(\text{PHENOL})$  (sidechain of tyrosine),  
 $\text{CH}_2\text{CH}_2\text{SCH}_3$  (sidechain of methionine),  
 $\text{CH}(\text{CH}_3\text{OH})$  (sidechain of threonine),  
 $\text{CH}_2\text{CONH}_2$  (sidechain of asparagine),  
10  $\text{CH}_2\text{CH}_2\text{CONH}_2$  (sidechain of glutamine),  
 $\text{CH}_2\text{CH}_2\text{COOH}$  (sidechain of glutamic acid),  
 $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_3^+$  (sidechain of lysine),  
 $\text{CH}_2\text{CH}_2\text{CH}_2\text{NC}(\text{NH}_2)_2$  (sidechain of arginine),  
 $\text{CH}_2(\text{IMIDAZOLE})$  (sidechain of histidine).
- 15 (2) There are no specific sequence requirements at positions  $P_4$ ,  $P_3$ ,  $P_2$ ,  $P_1$ ,  $P_2'$ ,  $P_3'$ , and  $P_4'$ . Each site can accommodate any other amino acid residue in singularity as long as rule number 3 is met.
- (3) At least two of the remaining seven positions,  $P_4$ ,  $P_3$ ,  $P_2$ ,  $P_1$ ,  $P_2'$ ,  $P_3'$ , and  $P_4'$ , must have an  $R_1$  which is made up of a hydrophobic residue. It is preferred that  
 20 there are at least three hydrophobic residues in the remaining seven positions,  $P_4$ ,  $P_3$ ,  $P_2$ ,  $P_1$ ,  $P_2'$ ,  $P_3'$ , and  $P_4'$ . Preferred  $R_1$  groups for the positions that contain a hydrophobic group are  $\text{CH}_3$  (side chain of alanine),  $\text{CH}(\text{CH}_3)_2$  (side chain of valine),  $\text{CH}_2\text{CH}(\text{CH}_3)_2$  (side chain of leucine),  $(\text{CH}_3)\text{CH}(\text{CH}_2\text{CH}_3)$  (side chain of isoleucine),  $\text{CH}_2(\text{indole})$  (side chain of tryptophan),  $\text{CH}_2(\text{benzene})$  (side chain of phenylalanine),  
 25  $\text{CH}_2\text{CH}_2\text{SCH}_3$  (side chain of methionine)  $\text{CH}_2(\text{phenol})$  (side chain of tyrosine). It is more preferred that the hydrophobic group be a large hydrophobic group. Preferred  $R_1$ s which contain large hydrophobic groups are  $\text{CH}(\text{CH}_3)_2$  (side chain of valine),  $\text{CH}_2\text{CH}(\text{CH}_3)_2$  (side chain of leucine),  $(\text{CH}_3)\text{CH}(\text{CH}_2\text{CH}_3)$  (side chain of isoleucine),  $\text{CH}_2(\text{indole})$  (side chain of tryptophan),  $\text{CH}_2(\text{benzene})$  (side chain of phenylalanine),  
 30  $\text{CH}_2\text{CH}_2\text{SCH}_3$  (side chain of methionine)  $\text{CH}_2(\text{phenol})$  (side chain of tyrosine). It is

-17-

most preferred that positions with a hydrophobic  $R_1$  are  $\text{CH}(\text{CH}_3)_2$  (side chain of valine),  $\text{CH}_2\text{CH}(\text{CH}_3)_2$  (side chain of leucine),  $\text{CH}_2(\text{benzene})$  (side chain of phenylalanine),  $\text{CH}_2\text{CH}_2\text{SCH}_3$  (side chain of methionine), or  $\text{CH}_2(\text{phenol})$  (side chain of tyrosine).

5 (4) None of the eight positions,  $P_4$ ,  $P_3$ ,  $P_2$ ,  $P_1$ ,  $P_1'$ ,  $P_2'$ ,  $P_3'$ , and  $P_4'$  may have a proline side chain at its  $R_1$  position.

(5) Not all subsites must have an P represented in the analog. For example, a substrate analog could have  $\text{X-P}_2\text{-L}_1\text{-P}_1\text{-L}_0\text{-P}_1'\text{-L}_1'\text{-P}_2'\text{-L}_2'\text{-P}_3'\text{-L}_3'\text{-Y}$  or it could have  $\text{X-L}_1\text{-P}_1\text{-L}_0\text{-P}_1'\text{-L}_1'\text{-P}_2'\text{-L}_2'\text{-P}_3'\text{-L}_3'\text{-P}_4'\text{-L}_4'\text{-Y}$ .

10 Preferred substrate analogs are analogs having the sequences disclosed in Table 1, with the non-hydrolyzable analog between  $P_1$  and  $P_1'$ .

#### Combinatorial Chemistry to Make Inhibitors

Combinatorial chemistry includes but is not limited to all methods for isolating molecules that are capable of binding either a small molecule or another  
 15 macromolecule. Proteins, oligonucleotides, and polysaccharides are examples of macromolecules. For example, oligonucleotide molecules with a given function, catalytic or ligand-binding, can be isolated from a complex mixture of random oligonucleotides in what has been referred to as "*in vitro* genetics" (Szostak, TIBS 19:89, 1992, the teachings of which are incorporated herein in their entirety). One  
 20 synthesizes a large pool of molecules bearing random and defined sequences and subjects that complex mixture, for example, approximately  $10^{15}$  individual sequences in 100  $\mu\text{g}$  of a 100 nucleotide RNA, to some selection and enrichment process. Through repeated cycles of affinity chromatography and PCR amplification of the molecules bound to the ligand on the column, Ellington and Szostak (1990)  
 25 estimated that 1 in  $10^{10}$  RNA molecules folded in such a way as to bind a small molecule dyes. DNA molecules with such ligand-binding behavior have been isolated as well (Ellington and Szostak, 1992; Bock et al, 1992, the teachings of all of which are incorporated herein in their entirety).

Techniques aimed at similar goals exist for small organic molecules, proteins  
 30 and peptides and other molecules known to those of skill in the art. Screening sets

-18-

of molecules for a desired activity whether based on libraries of small synthetic molecules, oligonucleotides, proteins or peptides is broadly referred to as combinatorial chemistry.

There are a number of methods for isolating proteins either have *de novo* activity or a modified activity. For example, phage display libraries have been used for a number of years. A preferred method for isolating proteins that have a given function is described by Roberts and Szostak (Roberts R.W. and Szostak J.W. Proc. Natl. Acad. Sci. USA, 94 (23):12997-302 (1997), the teachings of which are incorporated herein in their entirety). Another preferred method for combinatorial methods designed to isolate peptides is described in Cohen *et al.* (Cohen B.A., *et al.*, Proc. Natl. Acad. Sci. USA 95 (24):14272-7 (1998), the teachings of which are incorporated herein in their entirety). This method utilizes a modified two-hybrid technology. Yeast two-hybrid systems are useful for the detection and analysis of protein:protein interactions. The two-hybrid system, initially described in the yeast *Saccharomyces cerevisiae*, is a powerful molecular genetic technique for identifying new regulatory molecules, specific to the protein of interest (Fields and Song, *Nature* 340:245-6 (1989), the teachings of which are incorporated herein in their entirety). Cohen *et al.*, modified this technology so that novel interactions between synthetic or engineered peptide sequences could be identified which bind a molecule of choice. The benefit of this type of technology is that the selection is done in an intracellular environment. The method utilizes a library of peptide molecules that attach to an acidic activation domain. A peptide of choice, for example an extracellular portion of memapsin 2 is attached to a DNA binding domain of a transcriptional activation protein, such as Gal 4. By performing the Two-hybrid technique on this type of system, molecules that bind the extracellular portion of memapsin 2 can be identified.

#### *Screening of Small Molecule Libraries*

In addition to these more specialized techniques, methodology well known to those of skill in the art, in combination with various small molecule or combinatorial libraries, can be used to isolate and characterize those molecules which bind to or

-19-

interact with the desired target, either memapsin 2 or its substrate. The relative binding affinity of these compounds can be compared and optimum inhibitors identified using competitive or non-competitive binding studies which are well known to those of skill in the art. Preferred competitive inhibitors are non-  
5 hydrolyzable analogs of memapsin 2. Another will cause allosteric rearrangements which prevent memapsin 2 from functioning or folding correctly.

#### Computer assisted Rational Drug Design

Another way to isolate inhibitors is through rational design. This is achieved through structural information and computer modeling. Computer modeling  
10 technology allows visualization of the three-dimensional atomic structure of a selected molecule and the rational design of new compounds that will interact with the molecule. The three-dimensional construct typically depends on data from X-ray crystallographic analyses or NMR imaging of the selected molecule. The molecular dynamics require force field data. The computer graphics systems enable prediction  
15 of how a new compound will link to the target molecule and allow experimental manipulation of the structures of the compound and target molecule to perfect binding specificity. For example, using NMR spectroscopy, Inouye and coworkers were able to obtain the structural information of N-terminal truncated TSHK (transmembrane sensor histidine kinases) fragments which retain the structure of the  
20 individual sub-domains of the catalytic site of a TSHK. On the basis of the NMR study, they were able to identify potential TSHK inhibitors (U.S. Patent No. 6,077,682 to Inouye, the teachings of which are incorporated herein in their entirety). Another good example is based on the three-dimensional structure of a calcineurin/FKBP12/FK506 complex determined using high resolution X-ray  
25 crystallography to obtain the shape and structure of both the calcineurin active site binding pocket and the auxiliary FKBP12/FK506 binding pocket (U.S. Patent No. 5,978,740 to Armistead, the teachings of which are incorporated herein in their entirety). With this information in hand, researchers can have a good understanding of the association of natural ligands or substrates with the binding pockets of their

-20-

corresponding receptors or enzymes and are thus able to design and make effective inhibitors.

Prediction of molecule-compound interaction when small changes are made in one or both requires molecular mechanics software and computationally intensive computers, usually coupled with user-friendly, menu-driven interfaces between the molecular design program and the user. Examples of molecular modeling systems are the CHARMM and QUANTA programs, Polygen Corporation, Waltham, MA. CHARMM performs the energy minimization and molecular dynamics functions. QUANTA performs the construction, graphic modeling and analysis of molecular structure. QUANTA allows interactive construction, modification, visualization, and analysis of the behavior of molecules with each other.

A number of articles review computer modeling of drugs interactive with specific proteins, such as Rotivinen, *et al.*, 1988 *Acta Pharmaceutica Fennica* 97, 159-166; Ripka, *New Scientist* 54-57 (June 16, 1988); McKinaly and Rossmann, 1989 *Annu. Rev. Pharmacol. Toxicol.* 29, 111-122; Perry and Davies, QSAR: Quantitative Structure-Activity Relationships in Drug Design pp. 189-193 (Alan R. Liss, Inc. 1989; Lewis and Dean, 1989 *Proc. R. Soc. Lond.* 236, 125-140 and 141-162; and, with respect to a model enzyme for nucleic acid components, Askew, *et al.*, 1989 *J. Am. Chem. Soc.* 111, 1082-1090, the teachings of all of which are incorporated herein in their entirety). Other computer programs that screen and graphically depict chemicals are available from companies such as BioDesign, Inc., Pasadena, CA., Allelix, Inc, Mississauga, Ontario, Canada, and Hypercube, Inc., Cambridge, Ontario.

Although described above with reference to design and generation of compounds which could alter binding, one could also screen libraries of known compounds, including natural products or synthetic chemicals, and biologically active materials, including proteins, for compounds which alter substrate binding or enzymatic activity.

-21-

### Screening of Libraries

Design of substrate analogs and rational drug design are based on knowledge of the active site and target, and utilize computer software programs that create detailed structures of the enzyme and its substrate, as well as ways they interact, alone or in the presence of inhibitor. These techniques are significantly enhanced with X-ray crystallographic data in hand. Inhibitors can also be obtained by screening libraries of existing compounds for those which inhibit the catalytically active enzyme. In contrast to reports in the literature relating to memapsin 2, the enzyme described herein has activity analogous to the naturally produced enzyme, providing a means for identifying compounds which inhibit the endogenous activity. These potential inhibitors are typically identified using high throughput assays, in which enzyme, substrate (preferably a chromogenic substrate) and potential inhibitor (usually screened across a range of concentrations) are mixed and the extent of cleavage of substrate determined. Potentially useful inhibitors are those which decrease the amount of cleavage.

## **II. Preparation of Catalytically Active Recombinant Memapsin 2**

### Cloning and Expression of Memapsin 2

Memapsin 2 was cloned and the nucleotide (SEQ ID NO: 1) and predicted amino acid (SEQ ID NO: 2) sequences were determined, as described in Example 1. The cDNA was assembled from the fragments. The nucleotide and the deduced protein sequence are shown in SEQ ID NOS: 1 and 2, respectively. The protein is the same as the aspartic proteinase 2 (ASP2) described in EP 0 855 444 A by SmithKline Beecham Pharmaceuticals, (published July 29, 1998), U.S. Patent No. 6,319,689, Sinha, *et al.*, *Nature* 402, 537-540 (December 1999) and Vassar, *et al.*, *Science* 286, 735-741 (22 October 1999), the teachings of all of which are incorporated herein in their entirety).

Pro-memapsin 2 is homologous to other aspartic proteases, and shares homology with mouse ASP1 (U.S. Patent No. 6,291,223, the teachings of which are incorporated herein in its entirety). Based on the alignments, Pro-memapsin 2 contains a *pro* region, an aspartic protease region, and a trans-membrane region near

-22-

the C-terminus. The C-terminal domain is over 80 residues long. The active enzyme is memapsin 2 and its pro-enzyme is pro-memapsin 2.

#### Refolding Catalytically Active Enzyme

In order to determine the substrate specificity and to design inhibitors, it is  
5 necessary to express catalytically active recombinant enzyme. Since the active site is not in the transmembrane region and activity does not require membrane anchoring, memapsin 2 was expressed in *E. coli* in two different lengths, both without the transmembrane region, and purified, as described in Example 3. The procedures for the culture of transfected bacteria, induction of synthesis of recombinant proteins  
10 and the recovery and washing of inclusion bodies containing recombinant proteins are essentially as described by Lin *et al.*, (1994), the teachings of which are incorporated herein in its entirety. For refolding, the protein is dissolved in a strong denaturing/reducing solution such as 8 M urea/100 mM beta-mercaptoethanol. The rate at which the protein is refolded, and in what solution, is critical to activity. In  
15 one method, the protein is dissolved into 8 M urea/100 mM beta-mercaptoethanol then rapidly diluted into 20 volumes of 20 mM-Tris, pH 9.0, which is then slowly adjusted to pH 8 with 1 M HCl. The refolding solution is then kept at 4°C for 24 to 48 hours before proceeding with purification. In the second method, an equal volume of 20 mM Tris, 0.5 mM oxidized/1.25 mM reduced glutathione, pH 9.0 is  
20 added to rapidly stirred pro-memapsin 2 in 8 M urea/10 mM beta-mercaptoethanol. The process is repeated three more times with 1 hour intervals. The resulting solution is then dialyzed against sufficient volume of 20 mM Tris base so that the final urea concentration is 0.4 M. The pH of the solution is then slowly adjusted to 8.0 with 1 M HCl.  
25 The refolded protein is then further purified by column chromatography, based on molecular weight exclusion, and/or elution using a salt gradient, and analyzed by SDS-PAGE analysis under reduced and non-reduced conditions.

### III. Substrate Specificity and Enzyme Kinetics of Memapsin 2

The inhibitors can be screened for inhibition of binding and cleavage by memapsin 2 of its substrate.

#### Substrate Specificity

- 5 The presence of memapsin 2 (M2) in the brain indicated that it might hydrolyze the  $\beta$ -amyloid precursor protein (APP). As described below, detailed enzymatic and cellular studies demonstrated that M2 fits all the criteria of the  $\beta$ -secretase. The M2 three-dimensional structure modeled as a type I integral membrane protein. The model suggested that its globular protease unit can
- 10 hydrolyze a membrane anchored polypeptide at a distance range of 20-30 residues from the membrane surface. As a transmembrane protein of the brain, APP is a potential substrate and its beta-secretase site, located about 28 residues from the plasma membrane surface, is within in the range for M2 proteolysis.
- A synthetic peptide derived from this site (SEVKM/DAEFR) (SEQ ID NO:
- 15 5) was hydrolyzed by M2<sub>pd</sub> (modified M2 containing amino acids from Ala<sup>-8p</sup> to Ala<sup>326</sup>) at the beta-secretase site (marked by /). A second peptide (SEVNL/DAEFR) (SEQ ID NO: 6) derived from the APP beta-secretase site and containing the 'Swedish mutation' (Mullan, M. *et al.*, *Nature Genet.* 2:340-342 (1992), the teachings of which are incorporated herein in its entirety), known to elevate the level
- 20 of A $\beta$  production in cells (Citron, M. *et al.*, *Nature* 260:672-674 (1992), the teachings of which are incorporated herein in its entirety), was hydrolyzed by M2<sub>pd</sub> with much higher catalytic efficiency. Both substrates were optimally cleaved at pH 4.0. A peptide derived from the processing site of presenilin 1 (SVNM/AEGD) (SEQ ID NO: 7) was also cleaved by M2<sub>pd</sub> with less efficient kinetic parameters. A
- 25 peptide derived from the APP gamma-secretase site (KGGVVIATVIVK) (SEQ ID NO: 8) was not cleaved by M2<sub>pd</sub>. Pepstatin A inhibited M2<sub>pd</sub> poorly (IC<sub>50</sub> approximately approximately 0.3 mM). The kinetic parameters indicate that both presenilin 1 ( $k_{cat}$ , 0.67 s<sup>-1</sup>;  $K_m$ , 15.2 mM;  $k_{cat}/K_m$ , 43.8 s<sup>-1</sup>M<sup>-1</sup>) and native APP peptides ( $k_{cat}/K_m$ , 39.9 s<sup>-1</sup>M<sup>-1</sup>) are not as good substrates as the Swedish APP peptide
- 30 ( $k_{cat}$ , 2.45 s<sup>-1</sup>,  $K_m$ , 1 mM;  $k_{cat}/K_m$ , 2450 s<sup>-1</sup>M<sup>-1</sup>).



-24-

To determine if M2 possesses an APP beta-secretase function in mammalian cells, memapsin 2 was transiently expressed in HeLa cells (Lin, X., *et al.*, *FASEB J.* 7:1070-1080 (1993), the teachings of which are incorporated herein in its entirety), metabolically pulse-labeled with  $^{35}\text{S}$ -Met, then immunoprecipitated with anti-APP  
5 antibodies for visualization of APP-generated fragments after SDS-polyacrylamide electrophoresis and imaging. SDS-PAGE patterns of immuno-precipitated APP N $\beta$ -fragment (97 kD band) from the conditioned media (2 h) of pulse-chase experiments showed that APP was cleaved by M2. Controls transfected with APP alone and co-transfected with APP and M2 with Bafilomycin A1 added were performed. SDS-  
10 PAGE patterns of APP  $\beta$ C-fragment (12 kD) were immunoprecipitated from the conditioned media of the same experiment as discussed above. Controls transfected with APP alone; co-transfected with APP and M2; co-transfected with APP and M2 with Bafilomycin A1; transfections of Swedish APP; and co-transfections of Swedish APP and M2 were performed. SDS-PAGE gels were also run of immuno-  
15 precipitated M2 (70 kD), M2 transfected cells; untransfected HeLa cells after long time film exposure; and endogenous M2 from HEK 293 cells. SDS-PAGE patterns of APP fragments (100 kD betaN-fragment and 95 kD betaN-fragment) recovered from conditioned media after immuno-precipitation using antibodies specific for different APP regions indicated that memapsin 2 cleaved APP.

20 Cells expressing both APP and M2 produced the 97 kD APP beta N-fragment (from the N-terminus to the beta-secretase site) in the conditioned media and the 12 kD betaC-fragment (from the beta-secretase site to the C-terminus) in the cell lysate. Controls transfected with APP alone produced little detectable betaN-fragment and no beta C-fragment. Bafilomycin A1, which is known to raise the  
25 intra-vesicle pH of lysosomes/endosomes and has been shown to inhibit APP cleavage by beta-secretase (Knops, J. *et al.*, *J. Biol. Chem.* 270: 2419-2422 (1995), the teachings of which are incorporated herein in its entirety), abolished the production of both APP fragments beta N- and beta C- in co-transfected cells. Cells transfected with Swedish APP alone did not produce the beta C-fragment band in the  
30 cell lysate but the co-transfection of Swedish APP and M2 did. This Swedish beta C-fragment band is more intense than that of wild-type APP. A 97-kD beta N-band

-25-

is also seen in the conditioned media but is about equal intensity as the wild-type APP transfection.

These results indicate that M2 processes the beta-secretase site of APP in acidic compartments such as the endosomes. To establish the expression of transfected M2 gene, the pulse-labeled cells were lysed and immuno-precipitated by anti-M2 antibodies. A 70 kD M2 band was seen in cells transfected with M2 gene, which has the same mobility as the major band from HEK 293 cells known to express beta-secretase (Citron, M. *et al.*, *Nature* 260:672-674 (1992), the teachings of which are incorporated herein in its entirety). A very faint band of M2 is also seen, after a long film exposure, in untransfected HeLa cells, indicating a very low level of endogenous M2, which is insufficient to produce betaN- or betaC-fragments without M2 transfection. Antibody A $\beta_{1-17}$ , which specifically recognizes residues 1-17 in A $\beta$  peptide, was used to confirm the correct beta-secretase site cleavage. In cells transfected with APP and M2, both beta N- and beta C-fragments are visible using an antibody recognizing the N-terminal region of APP present in both fragments. Antibody A $\beta_{1-17}$  recognize the beta N-fragment produced by endogenous beta-secretase in the untransfected cells. This antibody was, however, unable to recognize the betaN-fragment known to be present in cells co-transfected with APP and M2. These observations confirmed that betaN-fragment is the product of beta-secretase site cut by M2, which abolished the recognition epitope of A $\beta_{1-17}$ .

In specificity studies, it was found that M2<sub>pd</sub> cleaved its *pro* peptide (2 sites) and the protease portion (2 sites) during a 16 h incubation after activation (Table 1). Besides the three peptides discussed above, M2<sub>pd</sub> also cleaved oxidized bovine insulin B chain and a synthetic peptide Nch. Native proteins were not cleaved by M2<sub>pd</sub>.

These same methods can be used in combination with the disclosed inhibitors to screen for efficacy in inhibiting the memapsin.

#### IV. Methods of diagnosis and treatment

Inhibitors can be used in the diagnosis and treatment and/or prevention of Alzheimer's disease and conditions associated therewith, such as elevated levels of

-26-

the forty-two amino acid peptide cleavage product, and the accumulation of the peptide in amyloid plaques.

#### Diagnostic Uses

5 The substrate analogs can be used as reagents for specifically binding to memapsin 2 or memapsin 2 analogs and for aiding in memapsin 2 isolation and purification or characterization, as described in the examples. The inhibitors and purified recombinant enzyme can be used in screens for those individuals more genetically prone to develop Alzheimer's disease.

#### Therapeutic Uses

10 Recombinant human memapsin 2 cleaves a substrate with the sequence LVNM/AEGD (SEQ ID NO: 9). This sequence is the *in vivo* processing site sequence of human presenilins. Both presenilin 1 and presenilin 2 are integral membrane proteins. They are processed by protease cleavage, which removes the N terminal sequence from the unprocessed form. Once processed, presenilin forms a  
15 two-chain heterodimer (Capell *et al.*, *J. Biol. Chem.* 273, 3205 (1998); Thinakaran *et al.*, *Neurobiol. Dis.* 4, 438 (1998); Yu *et al.*, *Neurosci Lett.* 2;254(3): 125-8 (1998), the teachings of all of which are incorporated herein in their entirety), which is stable relative to the unprocessed presenilins. Unprocessed presenilins are quickly degraded (Thinakaran *et al.*, *J. Biol. Chem.* 272, 28415 (1997); Steiner *et al.*, *J. Biol.*  
20 *Chem.* 273, 32322 (1998), the teachings of all of which are incorporated herein in their entirety). It is known that presenilin controls the *in vivo* activity of beta-secretase, which in turn cleaves the amyloid precursor protein (APP) leading to the formation of A $\beta$ 42. The accumulation of A $\beta$ 42 in the brain cells is known to be a major cause of Alzheimer's disease (for review, see Selkoe, 1998, the teachings of  
25 which are incorporated herein in its entirety). The activity of presenilin therefore enhances the progression of Alzheimer's disease. This is supported by the observation that in the absence of presenilin gene, the production of A $\beta$ 42 peptide is lowered (De Strooper *et al.*, *Nature* 391, 387 (1998), the teachings of which are incorporated herein in its entirety). Since unprocessed presenilin is degraded

-27-

quickly, the processed, heterodimeric presenilin must be responsible for the accumulation of A $\beta$ 42 leading to Alzheimer's disease. The processing of presenilin by memapsin 2 would enhance the production of A $\beta$ 42 and therefore, further the progress of Alzheimer's disease. Therefore a memapsin 2 inhibitor that crosses the blood brain barrier can be used to decrease the likelihood of developing or slow the progression of Alzheimer's disease which is mediated by deposition of A $\beta$ 42. Since memapsin 2 cleaves APP at the beta cleavage site, prevention of APP cleavage at the beta cleavage site will prevent the build up of A $\beta$ 42.

#### Pharmaceutically Acceptable Carriers

The inhibitors will typically be administered orally or by injection. Oral administration is preferred. Alternatively, other formulations can be used for delivery by pulmonary, mucosal or transdermal routes. The inhibitor will usually be administered in combination with a pharmaceutically acceptable carrier. Pharmaceutical carriers are known to those skilled in the art. The appropriate carrier will typically be selected based on the mode of administration. Pharmaceutical compositions may also include one or more active ingredients such as antimicrobial agents, antiinflammatory agents, and analgesics.

Preparations for parenteral administration or administration by injection include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Preferred parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, and electrolyte replenishers (such as those based on Ringer's dextrose).

Formulations for topical (including application to a mucosal surface, including the mouth, pulmonary, nasal, vaginal or rectal) administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Formulations for these applications are known. For example, a number of pulmonary formulations

-28-

have been developed, typically using spray drying to formulate a powder having particles with an aerodynamic diameter of between one and three microns, consisting of drug or drug in combination with polymer and/or surfactant.

Compositions for oral administration include powders or granules, suspensions or solutions in water or non-aqueous media, capsules, sachets, or tablets. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be desirable.

Peptides as described herein can also be administered as a pharmaceutically acceptable acid- or base- addition salt, formed by reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

#### Dosages

Dosing is dependent on severity and responsiveness of the condition to be treated, but will normally be one or more doses per day, with course of treatment lasting from several days to several months or until the attending physician determines no further benefit will be obtained. Persons of ordinary skill can determine optimum dosages, dosing methodologies and repetition rates.

The dosage ranges are those large enough to produce the desired effect in which the symptoms of the memapsin 2 mediated disorder are alleviated (typically characterized by a decrease in size and/or number of amyloid plaque, or by a failure to increase in size or quantity), or in which cleavage of the A $\beta$ 42 peptide is decreased. The dosage can be adjusted by the individual physician in the event of any counterindications.

The present invention will be further illustrated by the following examples which are not intended to be limiting in any way.

## EXEMPLIFICATION

**Example 1. Proteolytic activity and cleavage-site preferences of recombinant memapsin 2.**

The amino acid sequence around the proteolytic cleavage sites of APP was  
5 determined in order to establish the specificity of memapsin 2. Recombinant pro-  
memapsin 2-T1 was incubated in 0.1 M sodium acetate, pH 4.0, for 16 hours at  
room temperature in order to create autocatalyzed cleavages. The products were  
analyzed using SDS-polyacrylamide gel electrophoresis. Several bands which  
corresponded to molecular weights smaller than that of pro-memapsin 2 were  
10 observed. The electrophoretic bands were trans-blotted onto a PVDF membrane.  
Four bands were chosen and subjected to N-terminal sequence determination in a  
Protein Sequencer. The N-terminal sequence of these bands established the  
positions of proteolytic cleavage sites on pro-memapsin 2.

In addition, the oxidized  $\beta$ -chain of bovine insulin and two different  
15 synthetic peptides were used as substrates for memapsin 2 to determine the extent of  
other hydrolysis sites. These reactions were carried out by auto-activated pro-  
memapsin 2 in 0.1 M sodium acetate, pH 4.0, which was then incubated with the  
peptides. The hydrolytic products were subjected to HPLC on a reversed phase C-18  
column and the eluent peaks were subjected to electrospray mass spectrometry for  
20 the determination of the molecular weight of the fragments. Two hydrolytic sites  
were identified on oxidized insulin B-chain (Table1). Three hydrolytic sites were  
identified from peptide NCH-gamma. A single cleavage site was observed in  
synthetic peptide PS1-gamma, whose sequence (LVNMAEGD) (SEQ ID NO: 10) is  
derived from the beta-processing site of human presenilin 1 (Table 1).

-30-

**Table 1: Substrate Specificity of Memapsin 2**

Site #	Substrate	P4	P3	P2	P1	P1'	P2'	P3'	P4'	
1	Pro-memapsin 2	R	G	S	M	A	G	V	L	SEQ ID NO: 11 (aa 12-18 of SEQ ID NO: 3)
2		G	T	Q	H	G	I	R	L	SEQ ID NO: 12 (aa 23-30 of SEQ ID NO: 3)
3		S	S	N	F	A	V	G	A	SEQ ID NO: 13 (aa 98-105 of SEQ ID NO: 3)
4		G	L	A	Y	A	E	I	A	SEQ ID NO: 14 (aa 183-190 of SEQ ID NO: 3)
5	Oxidized Insulin B-Chain'	H	L	C^	G	S	H	L	V	SEQ ID NO: 15 C^ is cysteic acid;
6		C^	G	E	R	G	F	F	Y	SEQ ID NO: 16 C^ is cysteic acid
7	Synthetic Peptide*				V	G	S	G	V	Three sites cleaved in a peptide: VGSGVLLSRK(SEQ ID NO:30) SEQ ID NO: 17 SEQ ID NO: 18 SEQ ID NO: 19
8			V	G	S	G	V	L	L	
9		G	V	L	L	S	R	K		
10	Peptide**	L	V	N	M	A	E	G	D	SEQ ID NO: 10

Positions P<sub>n</sub> and P<sub>n</sub>' (wherein n is 1, 2, 3, 4, etc.) refer to the positioning of the amino acids in a peptide relative to the site of cleavage, indicated by the double vertical bar. Position numbers increase distally from the scissile bond.

\* Synthetic peptide based upon the amino acid sequence of Notch protein (Accession number AAG33848). Memapsin 2 cleaves the Notch peptide at three sites.

\*\* Synthetic peptide based upon the processing site of Presenilin-1 (Accession number P49768, Sherrington et al., Nature 375(6534): 754-760 (1995), the teachings of which are hereby incorporated by reference in its entirety).

**Example 2. Activation of pro-memapsin 2 and enzyme kinetics.**

Incubation in 0.1 M sodium acetate, pH 4.0, for 16 h at 22°C autocatalytically converted *pro*-M2<sub>pd</sub> to M2<sub>pd</sub>. For initial hydrolysis tests, two synthetic peptides were separately incubated with *pro*-M2<sub>pd</sub> in 0.1 M Na acetate, pH 4.0 for  
5 different periods ranging from 2 to 18 h. The incubated samples were subjected to LC/MS for the identification of the hydrolytic products. For kinetic studies, the identified HPLC (Beckman System Gold) product peaks were integrated for quantitation. The  $K_m$  and  $k_{cat}$  values for presenilin 1 and Swedish APP peptides (Table 1) were measured by steady-state kinetics. The individual  $K_m$  and  $k_{cat}$  values  
10 for APP peptide could not be measured accurately by standard methods, so its  $k_{cat}/K_m$  value was measured by competitive hydrolysis of mixed substrates against presenilin 1 peptide (Fersht, A. "Enzyme Structure and Mechanism", 2<sup>nd</sup> Ed., W.H. Freeman and Company, New York. (1985), the teachings of which are incorporated herein in their entirety).

15 The conversion of *pro*-M2<sub>pd</sub> at pH 4.0 to smaller fragments was shown by SDS-polyacrylamide electrophoresis. The difference in migration between *pro*-M2<sub>pd</sub> and converted enzyme is evident in a mixture of the two.

**Example 3: Design and Synthesis of Memapsin 2 Inhibitors OM99-1 and OM99-2.**

20 Based on the results of specificity studies of memapsin 2, it was predicted that good residues for positions P1 and P1' would be Leu and Ala. It was subsequently determined from the specificity data that P1' preferred small residues, such as Ala and Ser. However, the crystal structure (determined below) indicates that this site can accommodate a lot of larger residues. It was demonstrated that P1' of  
25 memapsin 2 is the position with the most stringent specificity requirement where residues of small side chains, such as Ala, Ser, and Asp, are preferred. Ala was selected for P1' mainly because its hydrophobicity over Ser and Asp is favored for the penetration of the blood-brain barrier, a requirement for the design of a memapsin 2 inhibitor drug for treating Alzheimer's disease. Therefore, inhibitors were designed  
30 to place a transition-state analogue isostere between Leu and Ala (shown as



-32-


Leu\*Ala, where \* represents the transition-state isostere,  $-\text{CH}(\text{OH})-\text{CH}_2-$  and the subsite P4, P3, P2, P2', P3' and P4' are filled with the beta-secretase site sequence of the Swedish mutant from the beta-amyloid protein.

OM99-1: Val-Asn-Leu\*Ala-Ala-Glu-Phe (SEQ ID NO: 20)

5 OM99-2: Glu-Val-Asn-Leu\*Ala-Ala-Glu-Phe (SEQ ID NO: 21)

The Leu\*Ala dipeptide isostere was synthesized as follows:

The Leu-Ala dipeptide isostere for the  $M_2$ -inhibitor was prepared from L-leucine. A bolded number in the following description, e.g. **2**, **3**, **4**, etc, refers to the respective numbered compound in synthesis schemes 1, 2, and 3. The bolded



(Scheme 1)

-33-


number is also referred to herein as "compound", e.g. compound 2, of the synthesis scheme.

As shown in Scheme 1, L-leucine was protected as its BOC-derivative 2 by treatment with  $\text{BOC}_2\text{O}$  in the presence of 10% NaOH in diethyl ether for 12 h. Boc-leucine 2 was then converted to Weinreb amide 3 by treatment with isobutyl chloroformate and N-methylpiperidine followed by treatment of the resulting mixed anhydride with N,O-dimethylhydroxylamine (Nahm and Weinreb, Tetrahedron Letters 1981, 32: 3815, the teachings of which are incorporated herein in their entirety). Reduction of 3 with lithium aluminum hydride in diethyl ether provided the aldehyde 4. Reaction of the aldehyde 4 with lithium propiolate derived from the treatment of ethyl propiolate and lithium diisopropylamide afforded the acetylenic alcohol 5 as an inseparable mixture of diastereomers (5.8:1) in 42% isolated yield (Fray, Kaye and Kleinman, J. Org. Chem. 1986, 51: 4828-33, the teachings of which are incorporated herein in their entirety). Catalytic hydrogenation of 5 over Pd/BaSO<sub>4</sub> followed by acid-catalyzed lactonization of the resulting gamma-hydroxy ester with a catalytic amount of acetic acid in toluene at reflux, furnished the gamma-lactone 6 and 7 in 73% yield.

The isomers were separated by silica gel chromatography by using 40% ethyl acetate in hexane as the eluent. Introduction of the methyl group at C-2 was accomplished by stereoselective alkylation of 7 with methyl iodide (Scheme 2). Thus, generation of the dianion of lactone 7 with lithium hexamethyldisilazide (2.2 equivalents) in tetrahydrofuran at -78°C (30 min) and alkylation with methyl iodide (1.1 equivalents) for 30 min at -78°C, followed by quenching with propionic acid (5 equivalents), provided the desired alkylated lactone 8 (76% yield) along with a small amount (less than 5%) of the corresponding epimer (Ghosh and Fidanze, 1998 J. Org. Chem. 1998, 63, 6146-54, the teachings of which are incorporated herein in their entirety). The epimeric cis-lactone was removed by column chromatography over silica gel using a mixture (3:1) of ethyl acetate and hexane as the solvent system. The stereochemical assignment of alkylated lactone 8 was made based on extensive <sup>1</sup>H-NMR NOE experiments. Aqueous lithium hydroxide promoted hydrolysis of the lactone 8 followed

-34-

by protection of the gamma-hydroxyl group with *tert*-butyldimethylsilyl chloride in the presence of imidazole and dimethylaminopyridine in dimethylformamide afforded the acid **9** in 90% yield after standard work-up and chromatography. Selective removal of the BOC-group was effected by treatment with trifluoroacetic



(Scheme 2)

- 5 acid in dichloromethane at 0°C for 1 h. The resulting amine salt was then reacted with commercial (Aldrich, Milwaukee) Fmoc-succinimide derivative in dioxane in the presence of aqueous NaHCO<sub>3</sub> to provide the Fmoc-protected L\*A isostere **10** in 65% yield after chromatography. Protected isostere **10** was utilized in the preparation of a random sequence inhibitor library.

## 10 Experimental procedure

### *N*-(*tert*-Butoxycarbonyl)-L-Leucine (compound **2**).

- To the suspension of 10 g (76.2 mmol) of L-leucine in 140 mL of diethyl ether was added 80 mL of 10 % NaOH. After all solid dissolves, 20 mL (87.1 mmol) of BOC<sub>2</sub>O was added to the reaction mixture. The resulting reaction mixture was stirred at 23°C for 12 h. After this period, the layers were separated and the aqueous layer was acidified to pH 1 by careful addition of 1 N aqueous HCl at 0°C. The resulting mixture was extracted with ethyl acetate (3 x 100 mL). The organic

-35-

layers were combined and washed with brine and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solvent was removed under reduced pressure to provide title product which was used directly for next reaction without further purification (yield, 97 %).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  4.89 (broad d, 1H,  $J = 8.3$  Hz), 4.31 (m, 1H), 1.74-1.49 (m, 3H),  
5 1.44 (s, 9H), 0.95 (d, 6H,  $J = 6.5$  Hz).

*N*-(*tert*-Butoxycarbonyl)-*L*-leucine-*N'*-methoxy-*N'*-methylamide (**compound 3**).

To a stirred solution of *N*,*O*-dimethylhydroxyamine hydrochloride (5.52 g, 56.6 mmol) in dry dichloromethane (25 mL) under  $\text{N}_2$  atmosphere at  $0^\circ\text{C}$ , -methylpiperidine (6.9 mL, 56.6 mmol) was added dropwise. The resulting mixture  
10 was stirred at  $0^\circ\text{C}$  for 30 min. In a separate flask, *N*-(*tert*-butoxycarbonyl)-*L*-leucine (**2**) (11.9 g, 51.4 mmol) was dissolved in a mixture of THF (45 mL) and dichloromethane (180 mL) under  $\text{N}_2$  atmosphere. The resulting solution was cooled to  $-20^\circ\text{C}$ . To this solution was added 1-methylpiperidine (6.9 mL, 56.6 mmol) followed by isobutyl chloroformate (7.3 mL, 56.6 mmol). The resulting mixture was  
15 stirred for 5 minutes at  $-20^\circ\text{C}$  and the above solution of *N*,*O*-dimethylhydroxyamine was added to it. The reaction mixture was kept  $-20^\circ\text{C}$  for 30 minutes and then warmed to  $23^\circ\text{C}$ . The reaction was quenched with water and the layers were separated. The aqueous layer was extracted with dichloromethane (3 x 100 mL). The combined organic layers were washed with 10% citric acid, saturated sodium  
20 bicarbonate, and brine. The organic layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated under the reduced pressure. The residue was purified by flash silica gel chromatography (25% ethyl acetate/hexane) to yield the title compound **3** (13.8 g, 97%) as a pale yellow oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  5.06 (broad d, 1H,  $J = 9.1$  Hz), 4.70 (m, 1H), 3.82 (s, 3H), 3.13 (s, 3H), 1.70 (m, 1H), 1.46-1.36 (m, 2H) 1.41  
25 (s, 9H), 0.93 (dd, 6H,  $J = 6.5, 14.2$  Hz).

*N*-(*tert*-Butoxycarbonyl)-*L*-leucinal (**compound 4**).

To a stirred suspension of lithium aluminum hydride (770 mg, 20.3 mmol) in dry diethyl ether (60 mL) at  $-40^\circ\text{C}$  under  $\text{N}_2$  atmosphere, was added *N*-*tert*-butoxycarbonyl-*L*-leucine-*N'*-methoxy-*N'*-methylamide (5.05 g, 18.4 mmol) in

-36-

diethyl ether (20 mL). The resulting reaction mixture was stirred for 30 min. After this period, the reaction was quenched with 10% NaHSO<sub>4</sub> solution (30 mL). The resulting reaction mixture was then warmed to 23°C and stirred at that temperature for 30 min. The resulting solution was filtered and the filter cake was washed by two  
5 portions of diethyl ether. The combined organic layers were washed with saturated sodium bicarbonate, brine and dried over anhydrous MgSO<sub>4</sub>. Evaporation of the solvent under reduced pressure afforded the title aldehyde **4** (3.41 g) as a pale yellow oil. The resulting aldehyde was used immediately without further purification. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.5 (s, 1H), 4.9 (s, 1H), 4.2 (broad m, 1H), 1.8-1.6 (m,  
10 2H), 1.44 (s, 9H), 1.49-1.39 (m, 1H), 0.96 (dd, 6H, J = 2.7, 6.5 Hz).

*Ethyl (4S,5S)-and (4R,5S)-5-[(tert-Butoxycarbonyl)amino]-4-hydroxy-7-methyloct-2-ynoate (compound 5).*

To a stirred solution of diisopropylamine (1.1 mL, 7.9 mmol) in dry THF (60 mL) at 0°C under N<sub>2</sub> atmosphere, was added n-BuLi (1.6 M in hexane, 4.95 mL, 7.9  
15 mmol) dropwise. The resulting solution was stirred at 0°C for 5 min and then warmed to 23°C and stirred for 15 min. The mixture was cooled to -78°C and ethyl propiolate (801 µL) in THF (2 mL) was added dropwise over a period of 5 min. The mixture was stirred for 30 min, after which N-Boc-L-leucinal **4** (1.55 g, 7.2 mmol) in 8 mL of dry THF was added. The resulting mixture was stirred at -78°C for 1 h.  
20 After this period, the reaction was quenched with acetic acid (5 mL) in THF (20 mL). The reaction mixture was warmed up to 23°C and brine solution was added. The layers were separated and the organic layer was washed with saturated sodium bicarbonate and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent under reduced pressure provided a residue which was purified by flash silica gel chromatography  
25 (15 % ethyl acetate / hexane) to afford a mixture (3:1) of acetylenic alcohols **5** (0.96 g, 42 %). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 4.64 (d, 1H, J = 9.0 Hz), 4.44 (broad s, 1H), 4.18 (m, 2H), 3.76 (m, 1H), 1.63 (m, 1H), 1.43-1.31 (m, 2H), 1.39 (s, 9H), 1.29-1.18 (m, 3H), 0.89 (m, 6H).

-37-

*(5S,1'S)-5-[1'-[(tert-Butoxycarbonyl)amino]-3'-methylbutyl]-dihydrofuran-2(3H)-one (compound 7).*

To a stirred solution of the above mixture of acetylenic alcohols (1.73 g, 5.5 mmol) in ethyl acetate (20 mL) was added 5% Pd/BaSO<sub>4</sub> (1 g). The resulting mixture was hydrogenated at 50 psi for 1.5 h. After this period, the catalyst was filtered off through a plug of Celite and the filtrate was concentrated under reduced pressure. The residue was dissolved in toluene (20 mL) and acetic acid (100  $\mu$ L). The reaction mixture was refluxed for 6 h. After this period, the reaction was cooled to 23°C and the solvent was evaporated to give a residue which was purified by flash silica gel chromatography (40% diethyl ether / hexane) to yield the (5S, 1S')-gamma-lactone 7 (0.94 g, 62.8% and the (5R, 1S')-gamma-lactone 6 (0.16 g, 10.7 %). Lactone 7: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  4.50-4.44 (m, 2H), 3.84-3.82 (m, 1H), 2.50 (t, 2H, J = 7.8 Hz), 2.22-2.10 (m, 2H), 1.64-1.31 (m, 3H), 1.41 (s, 9H), 0.91 (dd, 6H, J = 2.2, 6.7 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  177.2, 156.0, 82.5, 79.8, 51.0, 42.2, 28.6, 28.2, 24.7, 24.2, 23.0, 21.9.

*(3R,5S,1'S)-5-[1'-[(tert-Butoxycarbonyl)amino]]-3'-methylbut-yl]-3-methyl dihydrofuran-2(3H)-one (compound 8).*

To a stirred solution of the lactone 7 (451.8 mg, 1.67 mmol) in dry THF (8 mL) at -78°C under N<sub>2</sub> atmosphere, was added lithium hexamethyldisilazane (3.67 mL, 1.0 M in THF) over a period of 3 min. The resulting mixture was stirred at -78°C for 30 min to generate the lithium enolate. After this period, MeI (228  $\mu$ L) was added dropwise and the resulting mixture was stirred at -78°C for 20 min. The reaction was quenched with saturated aqueous NH<sub>4</sub>Cl solution and was allowed to warm to 23°C. The reaction mixture was concentrated under reduced pressure and the residue was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were washed with brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent afforded a residue which was purified by silica gel chromatography (15 % ethyl acetate / hexane) to furnish the alkylated lactone 8 (0.36 g, 76 %) as an amorphous solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  4.43 (broad t, 1H, J = 6.3 Hz), 4.33 (d, 1H, J = 9.6 Hz), 3.78 (m, 1H), 2.62 (m, 1H), 2.35 (m, 1H), 1.86 (m, 1H), 1.63-

-38-

1.24 (m, 3H), 1.37 (s, 9H), 1.21 (d, 3H, J = 7.5 Hz), 0.87 (dd, 6H, J = 2.6, 6.7 Hz);  
<sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 180.4, 156.0, 80.3, 79.8, 51.6, 41.9, 34.3, 32.5, 28.3,  
24.7, 23.0, 21.8, 16.6.

*(2R,4S,5S)-5-[(tert-Butoxycarbonyl)amino]-4-[(tert-butyldimethylsilyl)oxy]-2,7-*  
5 *dimethyloctanoic acid (compound 9)*

To a stirred solution of lactone 8 (0.33 g, 1.17 mmol) in THF (2 mL) was added 1 N aqueous LiOH solution (5.8 mL). The resulting mixture was stirred at 23°C for 10 h. After this period, the reaction mixture was concentrated under reduced pressure and the remaining aqueous residue was cooled to 0°C and acidified  
10 with 25% citric acid solution to pH 4. The resulting acidic solution was extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to yield the corresponding hydroxy acid (330 mg) as a white foam. This hydroxy acid was used directly for the next reaction without further purification.

15 To the above hydroxy acid (330 mg, 1.1 mmol) in anhydrous DMF was added imidazole (1.59 g, 23.34 mmol) and tert-butyldimethylchlorosilane (1.76 g, 11.67 mmol). The resulting mixture was stirred at 23°C for 24 h. After this period, MeOH (4 mL) was added and the mixture was stirred for 1 h. The mixture was diluted with 25% citric acid (20 mL) and was extracted with ethyl acetate (3 x 20  
20 mL). The combined extracts were washed with water, brine and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent gave a viscous oil which was purified by flash chromatography over silica gel (35% ethyl acetate / hexane) to afford the silyl protected acid 9 (0.44 g, 90 %). IR (neat) 3300-3000 (broad), 2955, 2932, 2859, 1711 cm<sup>-1</sup>; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>, 343 K) delta 6.20 (broad s, 1 H),  
25 3.68 (m, 1H), 3.51 (broad s, 1H), 2.49-2.42 (m, 1H), 1.83 (t, 1H, J = 10.1 Hz), 1.56 (m, 1H), 1.37 (s, 9H), 1.28-1.12 (m, 3H), 1.08 (d, 3H, J = 7.1 Hz), 0.87 (d, 3H, J = 6.1 Hz) 0.86 (s, 9 H), 0.82 (d, 3H, J = 6.5 Hz), 0.084 (s, 3H), 0.052 (s, 3H).

(2*R*,4*S*,5*S*)-5-[(fluorenylmethyloxycarbonyl)amino]-4-[(*tert*-butyldi-methylsilyl)oxy]-2,7-dimethyloctanoic acid (**compound 10**).

To a stirred solution of the acid 9 (0.17 g, 0.41 mmol) in dichloromethane (2 mL) at 0°C was added trifluoroacetic acid (500 µL). The resulting mixture was stirred at 0°C for 1 h and an additional portion (500 µL) of trifluoroacetic acid was added to the reaction mixture. The mixture was stirred for an additional 30 min and the progress of the reaction was monitored by TLC. After this period, the solvents were carefully removed under reduced pressure at a bath temperature not exceeding 5°C. The residue was dissolved in dioxane (3 mL) and NaHCO<sub>3</sub> (300 mg) in 5 mL of H<sub>2</sub>O. To this solution was added Fmoc-succinimide (166.5 mg, 0.49 mmol) in 5 mL of dioxane. The resulting mixture was stirred at 23°C for 8 h. The mixture was then diluted with H<sub>2</sub>O (5 mL) and acidified with 25% aqueous citric acid to pH 4. The acidic solution was extracted with ethyl acetate (3 x 50 mL). The combined extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure to give a viscous oil residue. Purification of the residue by flash chromatography over silica gel afforded the Fmoc-protected acid 10 (137 mg, 61%) as a white foam. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>, 343 K) δ 7.84 (d, 2H, J = 7.4 Hz), 7.66 (d, 2H, J = 8 Hz), 7.39 (t, 2H, J = 7.4 Hz), 7.29 (m, 2H), 6.8 (s, 1H), 4.29-4.19 (m, 3H), 3.74-3.59 (m, 2H), 2.49 (m, 1H), 1.88 (m, 1H), 1.58 (m, 1H), 1.31-1.17 (m, 3H), 1.10 (d, 3H, J = 7.1 Hz), 0.88 (s, 9H), 0.82 (d, 6H, J = 6.2 Hz), 0.089 (s, 3 H), 0.057 (s, 3H).

The synthesis of OM99-1 and OM99-2 were accomplished using solid state peptide synthesis procedure in which Leu\*Ala was incorporated in the fourth step. The synthesized inhibitors were purified by reverse phase HPLC and their structure confirmed by mass spectrometry.


#### Example 4: Design, Synthesis and Analysis of Additional Inhibitors.

Various substrate analogues other than OM99-1 and OM99-2 can be similarly designed and synthesized. For example, sixty six additional inhibitor analogues, MMI-001 to MMI-062, MMI-065, MMI-066, MMI-070 and MMI-071, all of which resemble an isostere of the active site of memapsin 2, were designed. The synthesis of the additional substrate analogues follows that for OM99-1 and



-40-

OM99-2. The chemical structures of the additional substrate analogues are listed in Table 3. The general synthesis of various inhibitors are outlined in Scheme 3.



Scheme 3: (a) EDC, HOBT, amine 11, *i*Pr<sub>2</sub>NEt, DMF-CH<sub>2</sub>Cl<sub>2</sub> (1:1), 0 °C to 23 °C;  
 (b) *n*Bu<sub>4</sub>N<sup>+</sup>F<sup>-</sup>, THF, 23 °C; (c) CF<sub>3</sub>CO<sub>2</sub>H, CH<sub>2</sub>Cl<sub>2</sub>, 23 °C;  
 (d) EDC, HOBT, acid, *i*Pr<sub>2</sub>NEt, DMF-CH<sub>2</sub>Cl<sub>2</sub> (1:1), 0 °C to 23 °C.

As scheme 3 demonstrates, using a standard peptide coupling procedure, valine derivative 11 was reacted with the dipeptide isostere 9 in the presence of N-ethyl-N'-(dimethylaminopropyl)carbodiimide hydrochloride, triethylamine and 1-hydroxybenzotriazole hydrate in a mixture of DMF and CH<sub>2</sub>Cl<sub>2</sub> to afford the amide derivative 12. Removal of the silyl protecting group by treatment with tetrabutylammonium fluoride in THF afforded inhibitor 13 (MMI-001). Exposure of 13 to trifluoroacetic acid in CH<sub>2</sub>Cl<sub>2</sub> resulted in the removal of BOC group. Coupling of the resulting amine with BOC-asparagine provided inhibitor 14 (MMI-011). Treatment of 14 with trifluoroacetic acid and coupling of the resulting amine with BOC-valine under standard conditions afforded inhibitor 16 (MMI-012). For the

-41-

synthesis of inhibitor MMI-15, compound 13 was reacted with trifluoroacetic acid and the resulting amine was coupled with BOC-methionine to afford the inhibitor 15 (MMI-015). Removal of the BOC group of 16 and coupling of the resulting amine with BOC-valine provided the inhibitor 17 (MMI-017).

5 ***Preparation of inhibitors MMI-001, MMI-011, MMI-012, MMI-015 and MMI-17:***

***Preparation of valine derivative (compound 11):*** N-Boc Valine (500 mg, 2.30 mmol) and benzylamine (0.50 mL, 4.60 mmol) were dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and DMF (2 mL). To this solution, HOBt (373 mg, 2.76 mmol) and EDC (529 mg, 2.76 mmol), and diisopropylethylamine (2.4 mL, 13.80 mmol) were added  
10 successively at 0°C. After the addition, the reaction mixture was allowed to warm up to 23°C and stirred overnight. The mixture was poured into sat. NaHCO<sub>3</sub> (aq). The resultant mixture was extracted with 30% EtOAc/hexane. The organic layer was washed with brine and dried over Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent under the reduced pressure gave a residue which was purified by flash column chromatography  
15 (30% EtOAc/hexane) to give 442 mg (63%) of coupling product. The resulting amine was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL). TFA (4 mL) was then added at room temperature. The reaction mixture was stirred for 0.5 hr. and it was concentrated under the reduced pressure. The amine 11 was obtained quantitative yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) 0.87 (3H, d, *J* = 6.9 Hz), 1.02 (3H, d, *J* = 6.9 Hz), 2.00 (2H, br s), 2.37 (1H, m), 3.36 (1H, br s), 4.43-4.52 (2H, m), 7.27-7.37 (5H, m), 7.70 (1H, br s).  
20 s).

***Preparation of amide derivative (compound 12):*** Dipeptide isostere 9 (41 mg, 0.10 mmol) and amine 11 (41 mg, 0.20 mmol) were dissolved in DMF (2.0 mL). To this solution, HOBt (20 mg, 0.15 mmol) and EDC (29 mg, 0.15 mmol),  
25 and diisopropylethylamine (0.2 mL) were added successively at 0°C. After the addition, the reaction mixture was allowed to warm up to 23°C and was stirred overnight. The mixture was poured into sat. NaHCO<sub>3</sub> (aq). The mixture was extracted with 30% EtOAc/hexane. The organic layer was washed with brine and

-42-

dried over  $\text{Na}_2\text{SO}_4$ . Evaporation of the solvent under the reduced pressure gave a residue which was purified by column chromatography (20% EtOAc/hexane) to give 55 mg (95%) of amide 12.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ) 0.09 (3H, s), 0.10 (3H, s), 0.91 (9H, s), 0.92-0.98 (12H, m), 1.10 (3H, d,  $J = 6.7$  Hz), 1.25 (1H, m), 1.44 (1H, m), 1.46 (9H, s), 1.63 (1H, m), 1.74 (1H, br s), 1.80 (1H, m), 2.18 (1H, m), 2.56 (1H, m), 3.62-3.78 (2H, m), 4.13 (1H, m), 4.48-4.56 (3H, m), 6.35 (1H, br d,  $J = 8.5$  Hz), 6.41 (1H, br s), 7.26-7.40 (5H, m).

**Preparation of inhibitor MMI-001 (compound 13):** To a solution of amide 12 (61 mg, 0.10 mmol) in THF (1.0 mL) was added TBAF (1.0 M in THF:0.3 mL, 0.30 mmol) at  $23^\circ\text{C}$  and it was stirred overnight. The reaction mixture was concentrated under reduced pressure. The residue was purified by column chromatography (40% EtOAc/hexane) to give MMI-001 (13, 41 mg, 83%).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) 0.88-0.98 (12H, m), 1.15 (3H, d,  $J = 6.9$  Hz), 1.40-1.80 (5H, m), 1.43 (9H, s), 2.10 (1H, m), 2.60 (1H, m), 3.40-3.60 (2H, m), 4.00 (1H, m), 4.20-4.45 (3H, m), 4.70 (1H, m), 7.08-7.18 (5H, m).

**Preparation of inhibitor MMI-011 (compound 14):** To a solution of 13 (44 mg, 0.089 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 mL) was added TFA (0.2 mL) at  $23^\circ\text{C}$ . After 0.5 hr, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF (4 mL). To this solution, *N*-Boc-asparagine (41 mg, 0.18 mmol), HOBt (24 mg, 0.18 mmol) and EDC (34 mg, 0.18 mmol), and diisopropylethylamine (0.2 mL) were added successively at  $0^\circ\text{C}$ . After the addition, the reaction mixture was allowed to warm up to  $23^\circ\text{C}$  and was stirred overnight. The mixture was poured into sat.  $\text{NaHCO}_3$  (aq). The mixture was extracted with EtOAc. The organic layer was washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation of the solvent under the reduced pressure gave a residue which was purified by column chromatography (4% MeOH/EtOAc) to give 12.5 mg (23%) of MMI-011 (14).  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}$ ) 0.85-1.00 (12H, m), 1.10 (3H, d,  $J = 6.7$  Hz), 1.20-1.50 (5H, m), 1.45 (9H, s), 1.60 (1H, m), 1.75 (1H, m), 2.05 (1H, m), 2.50-2.70 (3H, m), 3.45 (1H, m), 3.75 (1H, m), 4.10 (1H, m), 4.30-4.40 (3H, m), 7.20-7.30 (5H, m).

*Preparation of inhibitor MMI-015 (compound 15):* To a solution of compound 13 (64 mg, 0.13 mmol) in  $\text{CH}_2\text{Cl}_2$  (4 mL) was added TFA (1 ml) at 23°C. After 0.5 hr, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF (4 mL). To this solution, *N*-Boc-methionine (65 mg, 0.26 mmol), HOBt (35 mg, 0.26 mmol) and EDC (50 mg, 0.26 mmol), and diisopropylethylamine (0.4 mL) were added successively at 0°C. After the addition, the reaction mixture was allowed to warm up to 23°C and was stirred overnight. The mixture was poured into sat.  $\text{NaHCO}_3$  (aq). The mixture was extracted with EtOAc. The organic layer was washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation of the solvent under the reduced pressure gave a residue which was purified by column chromatography (80% EtOAc/hexane) to give 37.2 mg (46%) of MMI-015 (15).  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ) 0.80-1.00 (12H, m), 1.50 (3H, d,  $J = 6.7$  Hz), 1.20-2.10 (10H, m), 1.45 (9H, s), 2.10 (3H, s), 2.45-2.60 (2H, m), 2.70 (1H, m), 3.50 (1H, m), 3.90 (1H, m), 4.10-4.15 (2H, m), 4.30-4.42 (2H, m), 7.20-7.35 (5H, m).

*Preparation of Inhibitor MMI-012 (compound 16):* To a solution of 13 (8.4 mg, 0.014 mmol) in  $\text{CH}_2\text{Cl}_2$  (1 mL) was added TFA (0.2 ml) at 23°C. After 0.5 hr, the reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF (1 mL). To this solution, *N*-Boc-valine (6 mg, 0.028 mmol), HOBt (3.8 mg, 0.028 mmol) and EDC (5.3 mg, 0.028 mmol), and diisopropylethylamine (0.2 mL) were added successively at 0°C. After the addition, the reaction mixture was allowed to warm up to 23°C and was stirred overnight. The mixture was poured into sat.  $\text{NaHCO}_3$  (aq). The mixture was extracted with EtOAc. The organic layer was washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation of the solvent under reduced pressure gave a residue which was purified by column chromatography (4% MeOH/EtOAc) to give 2.1 mg (21%) of inhibitor MMI-012 (16).  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ) 0.80-1.00 (18H, m), 1.05 (3H, d,  $J = 6.7$  Hz), 1.10-1.50 (6H, m), 1.40 (9H, s), 1.55 (1H, m), 1.70 (1H, m), 2.00 (1H, m), 2.45-2.70 (3H, m), 3.45 (1H, m), 3.80-4.45 (6H, m), 7.15-7.30 (5H, m).

**Preparation of inhibitor MMI-107 (compound 17):** To a solution of 15 (14.5 mg, 0.023 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was added TFA (0.5 ml) at 23°C. After 0.5 hr. the reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF (2 mL). To this solution, *N*-Boc-valine (10 mg., 0.046 mmol), HOBt (6.2 mg, 0.046 mmol) and EDC (8.8 mg, 0.046 mmol), and diisopropylethylamine (0.4 mL) were added successively at 0°C. After the addition, the reaction mixture was allowed to warm up to 23°C and was stirred overnight. The mixture was poured into sat.  $\text{NaHCO}_3$  (aq). The mixture was extracted with EtOAc. The organic layer was washed with brine and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation of the solvent under the reduced pressure gave a residue which was purified by column chromatography (EtOAc) to give 5.5 mg (33%) of MMI-017 (17).  $R_f$  = (10% EtOAc/hexane);  $^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ) 0.80-1.00 (18H, m), 1.10 (3H, d,  $J$  = 6.7 Hz), 1.20-2.10 (11H, m), 1.43 (9H, s), 2.10 (3H, s), 2.40-2.60 (2H, m), 2.70 (1H, m), 3.50 (1H, m), 3.80-3.95 (2H, m), 4.18 (1H, m), 4.30-4.50 (3H, m), 7.20-7.30 (5H, m).

Enzyme activity of the isosteres was measured as described above, but with the addition of either OM99-1, OM99-2 or one of MMI-001 – MMI-062, MMI-065, MMI-066, MMI-070 and MMI-071. OM99-1 inhibited recombinant memapsin with a  $K_i$  calculated as  $3 \times 10^{-8}$  M. The substrate used was a synthetic fluorogenic peptide substrate. The inhibition of OM99-2 on recombinant memapsin 2 was measured using the same fluorogenic substrate. The  $K_i$  value was determined to be  $9.58 \times 10^{-9}$  M.

The residues in P1 and P1' are very important since the M2 inhibitor must penetrate the blood-brain barrier (BBB). The choice of Ala in P1' facilitates the penetration of BBB. Analogues of Ala side chains will also work. For example, in addition to the methyl side chain of Ala, substituted methyl groups and groups about the same size like methyl or ethyl groups can be substituted for the Ala side chain. Leu at P1 can also be substituted by groups of similar sizes or with substitutions on Leu side chain. For penetrating the BBB, it is desirable to make the inhibitors smaller. One can therefore use OM99-1 as a starting point and discard the outside subsites P4, P3, P3' and P4'. The retained structure Asn-Leu\*Ala-Ala is then further

-45-

evolved with substitutions for a tight-binding M2 inhibitor which can also penetrate the BBB.

The other substrate analogues, MMI-001 to MMI-071, were also tested for enzyme inhibition. For example, the  $K_i$  values for MMI-017, MMI-070 and MMI-071 are comparable to that for OM99-2, indicating that they are also excellent memapsin inhibitors. The  $K_i$  value for MMI-012, MMI-018, MMI-026, or MMI-066 is approximately one magnitude higher than that for OM99-2, indicating that they are competent candidates for memapsin inhibition. The  $K_i$  value for each of the additional substrate analogues and its corresponding chemical structure are listed in Table 3.

Table 3. Chemical Structures and  $K_i$  Values of Additional Substrate Analogues





















Structure	Compd No. / $K_i$ (nM)	Structure	Compd No./ $K_i$ (nM)
	MMI-001 / 3,738,100.0		MMI-011 / 3,134.0
	MMI-002 / 430,940.0		MMI-012 / 5.9
	MMI-003 / 2,617,000.0		MMI-013 / 184,160.0
	MMI-004 / 22,423.0		MMI-014 / 2,777.8
	MMI-005 / 61.4		MMI-015 / 5,808.0
	MMI-006 / 63,288.0		MMI-016 / 5,640.0
	MMI-007 / 49,877.0		MMI-017 / 2.5
	MMI-008 / 73.6		MMI-018 / 8.0
	MMI-009 / 89.9		MMI-019 / 14,500.0
	MMI-010 / 29,403.0		MMI-020 / 1,605.2

Table 3 cont'd from previous page





















	MMI-021 / 112.2		MMI-031 / 863.7
	MMI-022 / 11,766.0		MMI-032 / 1,341.0
	MMI-023 / 15,273.0		MMI-033 / 16,300.0
	MMI-024 / 1129.0		MMI-034 / 228.8
	MMI-025 / 50.1		MMI-035 / 426.3
	MMI-026 / 9.4		MMI-036 / no inhibition
	MMI-027 / 11,547.3		MMI-037 / 3.9
	MMI-028 / 9,867.0		MMI-038 / 106.0
	MMI-029 / 1,696.6		MMI-039 / 1,4
	MMI-030 / 8,517.0		MMI-040 / 2.1



Table 3 cont'd from previous page



























	MMI-041 / no inhibition		MMI-051 / no inhibition (polar isomer)
	MMI-042 / 36,484.1		MMI-052 / 5,790.0 (polar isomer)
	MMI-043 / 49,188.9		MMI-053 / no inhibition
	MMI-044 / no inhibition		MMI-054 / 469,000.0
	MMI-045 / no inhibition		MMI-055 / 631,000.0 (less polar isomer)
	MMI-046 / 149,900,000.0		MMI-056 / 205,000.0 (less polar isomer)
	MMI-047 / 110,600,000.0		MMI-057 / 73,300.0 (polar isomer)
	MMI-048 / 54,8000,000.0		MMI-058 / 408,000.0 (polar isomer)
	MMI-049 / 198,000.0		MMI-059 / 66,300.0 (less polar isomer)
	MMI-050 / 146,000.0		MMI-060 / 100,000.0

Table 3 cont'd from previous page

	MMI-061 / 15,460,000.0		MMI-066 / 21.2
	MMI-062 / no inhibition		MMI-070 / 2.0 (4.9)
	MMI-065 / 40.0 (97.4)		MMI-071 / 4.0 (1.9)

**Example 5. Sub-site Specificity of Memapsin 2 ( $\beta$ -secretase)**

The results obtained from OM99-2 demonstrate that by evoking all eight sub-sites with OM99-2, a high inhibition potency ( $K_i = 1.6$  nM) can be achieved.

However, clinically useful memapsin 2 inhibitors must be small (i.e., typically less than 500 - 700 daltons) in order to penetrate the blood-brain barrier to reach the brain. Detailed information on sub-site specificity of memapsin 2 would provide a better understanding for the design of small yet tight-binding inhibitors. The complete sub-site preference of memapsin 2 has been determined based on results from both substrate kinetics and the probing of random sequence inhibitor library.

**Experimental Procedures*****The Design of the Defined Substrate Mixtures***

Peptide mixtures were synthesized to probe each of 8 positions in the template sequence EVNLAAEF (SEQ ID NO: 22), derived from the  $\beta$ -secretase cleavage site in APP and memapsin 2 inhibitor OM99-2 described above. For characterization of each position, an equimolar mixture of 7 amino acid derivatives were added in the appropriate cycle of solid phase synthesis (Research Genetics, Invitrogen, Huntsville, Alabama) resulting in a mixture of 7 peptides, differing by 1 of 7 amino acids at a single position. To facilitate the rapid quantitation of substrates

-50-

and hydrolytic products, high throughput MALDI-TOF MS was used. Therefore, it was not possible to substitute all common amino acid side chains (19 total, excluding cysteine) in a single mixture due to the identical masses of some amino acids which would prevent their identification. Therefore, the different amino acid substitutions were divided into three groups (Tables 4 and 5) to maximize their differences in mass and the full set of 19 varied amino acids (less cysteine) at each subsite were contained in 3 substrate mixtures in which the peptides were synthesized to incorporate each amino acid in a single position. A standard peptide was included in each group for the purpose of relating the quantitative information between mixtures, for calculation of relative  $k_{cat}/K_m$ . The substrate of known  $k_{cat}/K_m$  value serves as an internal standard for normalization of relative initial rates and the calculation of  $k_{cat}/K_m$  value of other substrates. For positions  $P_1'$ ,  $P_2'$ ,  $P_3'$ , and  $P_4'$ , the template sequence was extended by 4 residues at the C-terminus (template EVNLAAEFWHDR, SEQ ID NO: 23, Table 4) to facilitate their detection in MALDI-TOF MS. Four additional residues were likewise added on the N-terminus in the template to characterize specificity for positions  $P_1$ ,  $P_2$ ,  $P_3$ , and  $P_4$  (template RWHHEVNLAEEF, SEQ ID NO: 24, Table 5).

**Table 4. Peptide Substrate Mixture for Specificity Determination of Memapsin 2 to the P<sub>1</sub>' Position**

	Amino Acid	Peptide Sequence	SEQ ID NO.
Substrate mixture 1	<b>A (alanine)</b>	EVNLAAEFWHDR	25
	D (aspartic acid)	EVNLDAEFWHDR	26
	S (serine)	EVNLSAEFWHDR	27
	T (threonine)	EVNLTAEFWHDR	28
	I (isoleucine)	EVNLIAEFWHDR	29
	E (glutamic acid)	EVNLEAEFWHDR	30
	F (phenylalanine)	EVNLF AEFWHDR	31
Substrate mixture 2	<b>A (alanine)</b>	EVNLAAEFWHDR	32
	G (glycine)	EVNLGAEFWHDR	33
	V (valine)	EVNLVAEFWHDR	34
	L (leucine)	EVNLLAEFWHDR	35
	K (lysine)	EVNLKAEFWHDR	36
	R (arginine)	EVNLR AEFWHDR	37
	W (tryptophan)	EVNLWAEFWHDR	38
Substrate mixture 3	<b>A (alanine)</b>	EVNLAAEFWHDR	39
	P (proline)	EVNLP AEFWHDR	40
	N (asparagine)	EVNLN AEFWHDR	41
	Q (glutamine)	EVNLQ AEFWHDR	42
	M (methionine)	EVNLM AEFWHDR	43
	H (histidine)	EVNLH AEFWHDR	44
	Y (tyrosine)	EVNLY AEFWHDR	45

**Table 5. Peptide Substrate Mixture for Specificity Determination of Memapsin 2 to the P<sub>1</sub> Position**

	Amino Acid	Peptide Sequence	SEQ ID NO.
Substrate mixture 1	<b>F (phenylalanine)</b>	RWHHEVNFAAEF	46
	D (aspartic acid)	RWHHEVNDAAEF	47
	S (serine)	RWHHEVNSAAEF	48
	T (threonine)	RWHHEVNTAAEF	49
	I (isoleucine)	RWHHEVNIAAEF	50
	E (glutamic acid)	RWHHEVNEAAEF	51
	G (glycine)	RWHHEVNGAAEF	52
5 Substrate mixture 2	<b>F (phenylalanine)</b>	RWHHEVNFAAEF	53
	A (alanine)	RWHHEVNAAAEF	54
	V (valine)	RWHHEVNVAAEF	55
	L (leucine)	RWHHEVNLAAEF	56
	Q (glutamine)	RWHHEVNQAAEF	57
	M (methionine)	RWHHEVNMAAEF	58
	Y (tyrosine)	RWHHEVNYAAEF	59
Substrate mixture 3	<b>F (phenylalanine)</b>	RWHHEVNFAAEF	60
	P (proline)	RWHHEVNPAAEF	61
	N (asparagine)	RWHHEVNNAAEF	62
	K (lysine)	RWHHEVNKAAEF	63
	R (arginine)	RWHHEVNRAAEF	64
	H (histidine)	RWHHEVNHAAEF	65
	W (tryptophan)	RWHHEVNWAAEF	66

***Initial Rate Determination by MALDI-TOF MS***

Substrate mixtures were dissolved at 2 mg/ml in 10% glacial acetic acid, and diluted into appropriate concentration of NaOH to obtain a mixture of substrates in the  $\mu$ M range in sodium acetate at pH 4.1. Aliquots were equilibrated at 25 °C, and reactions were initiated by the addition of aliquots of memapsin 2. Aliquots (10  $\mu$ l) were removed at time intervals and combined with MALDI-TOF matrix ( $\alpha$ -hydroxycinnamic acid in acetone, 20 mg/ml) and immediately spotted in duplicate onto a stainless-steel MALDI sample plate. Samples were subjected to analysis by using MALDI-TOF mass spectrometry device, operated at 20,000 accelerating volts in positive mode with a 150 nsec delay, using a PE Biosystems Voyager DE instrument at the Molecular Biology Resource Center on campus. Ions with a mass-to-charge ratio ( $m/z$ ) were detected in the range of 400 - 2000 amu (atomic mass

-53-

units). Data were analyzed by the Voyager Data Explorer module to obtain ion intensity data for mass species of interest.

### ***Random Sequence Inhibitor Library***

The random sequence inhibitor library was based on the sequence of OM99-2 with random amino acids (less cysteine) at 4 subsite positions  $P_2$ ,  $P_3$ ,  $P_2'$  and  $P_3'$ . Diisostere Leu\*Ala (\* represents hydroxyethylene transition-state isostere) was used in the synthesis to fix the positions  $P_1$  and  $P_1'$ . Peptides were synthesized by solid-state peptide synthesis method and left attached on the resin beads. By using the 'split-synthesis' procedure (Lam *et al.*, 1991), each of the resin beads contained only one sequence while the sequence differed from bead to bead. The overall library sequence was:

Gly-Xx1-Xx2-Leu\*Ala-Xx3-Xx4-Phe-Arg-Met-Gly-Gly-[Resin bead]

$P_4 \quad P_3 \quad P_2 \quad P_1 \quad P_1' \quad P_2' \quad P_3' \quad P_4'$

(SEQ ID NO: 67)

where Xxn residues (where n represents either 1, 2, 3, or 4) are randomized at each position with 19 amino acids. A shorter version of the peptides, starting at  $P_2'$  (sequence: Xx3-Xx4-Phe-Arg-Met-Gly-Gly-[Resin bead] SEQ ID NO: 68), was also present in each bead with a ratio to the longer sequence at about 7:3. Without isostere, the short sequence would not bind memapsin 2 with significant strength but its presence was convenient for identifying the residues at by automated Edman degradation. The residues are identified from the randomized positions as follows:

Edman cycle #	1	2	3	4
Sequence 1, from the long sequence:	Gly	Xx1	Xx2	Leu
Sequence 2, from the short sequence:	Xx3	Xx4	Phe	Arg

The assignment of Xx3 and Xx2 had no ambiguity since they are the only unknown residue at cycle 1 and 3 respectively. Amino acids Xx1 and Xx4 were assigned from their relative amounts. The presence of a methionine would permit the CNBr cleavage follow by peptide identified by MS/MS.

***Probing of the Random Sequence Library***

About 130,000 individual beads representing one copy of the library, estimated to be contained in 1.1 ml of settled beads, was hydrated in buffer A (50 mM Na acetate, 0.1% Triton X-100, 0.4 M urea, 0.02% Na azide, 1 mg/ml BSA, pH 3.5; filtered with a 5 micron filter). The beads were soaked in 3% BSA in buffer A for 1 h to block the non-specific binding, and rinsed twice with the same buffer. Recombinant memapsin 2 was diluted into buffer A to 4 nM and incubated with the library for 1 h. A single stringency wash was performed which included 6.7  $\mu$ M transition-state isosteric inhibitor OM99-2 in buffer B (50 mM Na acetate, 0.1% Triton X-100, 0.02% Na azide, 1 mg/ml BSA, pH 5.5; filtered with 5 micron filter), followed by two additional washes with buffer B containing no OM99-2. Affinity-purified IgG specific for recombinant memapsin 2 was diluted 100 fold in buffer B and incubated 30 min with the library. Following three washes with buffer B, affinity-purified anti-goat-alkaline phosphatase conjugate was diluted into buffer B (1:200) and incubated for 30 min, with three subsequent washes. A single tablet of alkaline phosphatase substrate (BCIP/NBT) was dissolved in 10 ml water and 1 ml applied to the beads and incubated 1 h. Beads were resuspended in 0.02% sodium azide in water and examined under a dissecting microscope. Darkly-stained beads were graded by sight, individually isolated, stripped in 8 M urea for 24 h, and destained in DMF. The sequence determination of the beads were carried out in an Applied Biosystem Protein Sequencer at the Molecular Biology Resource Center on campus and the PTH-amino acids were quantified using reversed-phase HPLC.

***Determination of Kinetic Parameters***

The kinetic parameters,  $K_m$  and  $k_{cat}$ , using single peptide substrate, and  $K_i$  against free inhibitors, were determined as described above.

***Model building of OM00-3 binding to memapsin 2***

The crystal structure of OM99-2 in complex with memapsin 2 was used as the initial model. Corresponding subsite residues were replaced with that of OM00-3. The side chain conformations were visually selected from a rotamer library for the best fitting into the binding pockets in terms of the non-bonded interaction.

-55-

Energy minimization was then carried out with the CNS program using Engh and Huber energy parameters (Engh, R.A. and Huber "Accurate bond and angle parameters for X-ray structure refinement" in: Acta Cryst. A47, 392-400 (1991)). C<sub>α</sub> atoms and the atoms with a distance of more than 5 angstroms from the inhibitor were subject to harmonic restrains during the minimization process.

### Results

#### *Determination of substrate side-chain preference in memapsin 2 sub-sites*

The substrate cleft of memapsin 2 accommodates eight sub-sites for the side chains as shown in the crystal structure. A complete set of side-chain preference analyzed by classical enzyme kinetics for all sub-sites of memapsin 2 would require the determination of 160 pairs of individual  $k_{cat}$  and  $K_m$  values, a tedious task so far not attained for any aspartic protease with broad specificity. The sub-site preference is, however, defined by the relative catalytic efficiency,  $k_{cat}/K_m$  values of substrate with different side chains, which may be determined from the relative initial hydrolysis rates of defined mixtures of substrates (Fersht, A. Enzyme Structure and Mechanism, 2<sup>nd</sup> Ed., Freeman, New York, (1985), the teachings of which are incorporated herein in their entirety). This method has been successfully used to analyze the specificity of other aspartic proteases (Kassel *et al.*, 1995; Koelsch *et al.*, 2000, the teachings of which are incorporated herein in their entirety). The rate determination was further simplified by the use of MALDI-TOF/MS ion intensities for quantification of relative amounts of products and substrates. MALDI-TOF/MS ion intensities have been used to quantify compounds from plasma and cell culture (Sugiyama *et al.*, "A quantitative analysis of serum sulfatide by matrix-assisted laser desorption ionization time-of-flight mass spectrometry with delayed ion extraction" in: *Anal. Biochem.* 274: 90-97 (1999); Wu *et al.*, "An automated MALDI mass spectrometry approach for optimizing cyclosporin extraction and quantitation" in: *Anal Chem.* 69: 3767-71 (1997), the teachings of all of which are incorporated herein in their entirety) and pertinently, to determine relative amount of A $\beta$  40 and A $\beta$  42 from cell culture (Davies *et al.*, "The structure and function of the aspartic proteinases" in: *Annu Rev Biophys Biophys Chem.* 19: 189-215 (1990), the teachings of which are incorporated herein in their entirety). Advantages of MALDI-TOF/MS



-56-

for this method include its sensitivity and rapid acquisition of data. Linearity of ion intensity data with mixtures of product and substrate produced excellent correlation, was consistent for each substrate in the mixture, and required no correction factor. Initial rates of hydrolysis for each peptide in each mixture were subsequently  
5 determined using this method. Ratios of these initial rates are proportional to their relative catalytic efficiencies,  $k_{cat}/K_m$  (Fersht, 1985).

The relative catalytic efficiencies of memapsin 2 for eight sub-site positions of the substrate are presented in Figure 5. These results confirm previous observations that none of the eight sub-site of memapsin 2 is absolutely stringent.  
10 On both the P side and the P' side, the central sub-sites ( $P_1$  and  $P_1'$ ) are more stringent than the outside sub-sites ( $P_4$  and  $P_4'$ ). This is in evidence when the hydrolysis efficiency of the non-preferred residues (background) are compared to the preferred residues. The lack of stringency is more pronounced for the 4 sub-sites on the P' side, especially for  $P_3'$  and  $P_4'$  where the backgrounds are relatively high. The  
15 poor stringencies of these two sub-sites is consistent with the observation that the  $P_3'$  and  $P_4'$  side chains of OM99-2 do not significantly interact with the enzyme in the the crystal structure.

Sub-site  $P_4$  favors Glu over Gln which, in turn, is favored over Asp.  $P_4$ -Glu of OM99-2 fits well in S4 pocket with multiple interactions. The reduction of  
20 catalytic efficiency from substitution of  $P_4$ -Glu with Gln may be due to the loss of charge interaction of  $P_4$  side chain with Arg<sup>235</sup> and Arg<sup>307</sup>. The further decrease of catalytic efficiency from substitution of  $P_4$ -Glu with Asp is likely due to the absence of the hydrogen bond to  $P_2$ -Asn. For sub-site  $P_3$ , an Ile is more preferable than Val as in OM99-2. This is due to a better fitting in the  $S_3$  hydrophobic pocket.

## 25 *Inhibitor side chain preference determined from a combinatorial Library*

The sub-site preference for inhibitor binding was also studied. A combinatorial library of approximately  $1.3 \times 10^5$  different inhibitors immobilized on beads was synthesized and probed with memapsin 2. The base sequence of the library was taken from OM99-2, EVNL\*AAEF (SEQ ID NO: 69) (\* designates  
30 isostere hydroxyethylene), in which the sub-sites  $P_3$ ,  $P_2$ ,  $P_2'$ , and  $P_3'$  (boldface) were randomized with all amino acids except cysteine.  $P_1$  and  $P_1'$  were keep constant due

-57-

to the use of L\*A in library synthesis. P4 and P4' were not randomized in order to keep the library size manageable. Also, the information on those outside sub-sites are less critical for the design of smaller inhibitors. After enhancing memapsin 2 binding selectivity to the library by washing with urea solutions, the bound

5 memapsin 2 on beads was detected by antibody raised to memapsin 2 and an alkaline phosphatase conjugated secondary antibody. About 65 from about 130,000 beads were intensely stained as observed under the light microscope. The inhibitor sequences from the 10 most intensely stained beads were determined by automated Edman sequencing (Table 2). A clear consensus at the four randomized positions

10 (boldface) was observed to be ELDLAVEF (SEQ ID NO: 70). The clear consensus in positive beads and the lack of consensus in negative beads (Table 2) supports the contention that the memapsin 2 was bound to preferred sequences in the library. Also, the consensus inhibitor sequence agree well with the results obtained from the substrate studies. An inhibitor, OM00-3, having the consensus sequence

15 ELDL\*AVEF (SEQ ID NO: 71) was synthesized using the method described above. The  $K_i$  of this inhibitor proved to be 0.31 nM, nearly five fold lower than the  $K_i$  of OM99-2.

Table 2. Selection of potent competitors of OM99-2 from combinatorial library<sup>a</sup>

ID	P3	P2	P2'	P3'	SEQ ID NO.
1	Leu	Asp	Val	Glu	72
2	Leu	Glu	Val	Glu	73
3	Leu	Asp	Val	Glu	74
4	Leu	Asp	Val	Glu	75
5	Leu	Asp	Val	Gln	76
6	Ile	Asp	Ala	Gln	77
7	Ile	Asp	Val	Tyr	78
8	Leu	Glu	Val	Gln	79
9	Leu	Phe	Val	Glu	80
10	Phe/Ile	Ser	Val	Phe/Ile	81
Neg1	Phe	Met	Asn	Arg	82
Neg2	Asp	Phe		Ser	83

<sup>a</sup>Library template: Gly-P<sub>3</sub>-P<sub>2</sub>-Leu\*Ala-P<sub>2</sub>'-P<sub>3</sub>'-Phe-Arg-Met-Gly-Gly-Resin  
(SEQ ID NOS: 72-83)

#### ***Binding of OM00-3 in the active site of memapsin 2***

Comparisons between the crystal structure of OM99-2 and the molecular model of OM2000-1 reveal improved inhibitor binding of the later to memapsin 2. As compared to OM99-2, the larger hydrophobic side chains at P<sub>3</sub> and P<sub>2</sub>' subsites for OM00-3 are better accommodated by the enzyme subsite through improved van der Waal interactions. Among these are six significant new interactions: P<sub>3</sub>-Leu of inhibitor with Leu and Gly of the enzyme and P<sub>2</sub>'-Val with Val, Tyr, Ile and Tyr. In addition, the model shows that replacement of Asn by Asp at the P<sub>2</sub> position introduces three new salt bridges between the side chain atoms CD1 and CD2 of P<sub>2</sub>-Asp and that of NE and NH1 of Arg-235. The inter-atomic distances of these salt bridges are 3.6, 3.4 and 3.5 angstroms respectively. These charge interactions

-59-

should render higher free energies of binding than that of P<sub>2</sub>-Asn in OM99-2. These observed structural differences between the binding of two inhibitors are consistent with the 4.5 fold decrease in K<sub>i</sub> values from OM99-2 to OM00-3.

To be useful for clinical treatment of AD, memapsin 2 inhibitors must  
5 penetrate the blood-brain barrier, and consequently, must be small in size (< 700 daltons). An eight-residue inhibitor such as OM00-3, is 917 daltons. Smaller inhibitors can be designed using fewer sub-sites, perhaps spanning only 4 or 5 sub-sites. Information described herein on sub-site preferences together with the information on the three-dimensional structure of the sub-sites can be used in the  
10 design of these inhibitors.

#### **Example 6: Using The Crystal Structure to Design Inhibitors**

Pharmaceutically acceptable inhibitor drugs are normally less than 800 daltons. In the case of memapsin 2 inhibitors, this requirement may even be more stringent due to the need for the drugs to penetrate the blood-brain. In the current  
15 model, well defined subsite structures at P<sub>4</sub> to P<sub>2</sub>' provide sufficient template areas for rational design of such drugs. The spacial relationships of individual inhibitor side chain with the corresponding subsite of the enzyme as revealed in this crystal structure permits the design of new inhibitor structures in each of these positions. It is also possible to incorporate the unique conformation of subsites P<sub>2</sub>', P<sub>3</sub>' and P<sub>4</sub>'  
20 into the selectivity of memapsin 2 inhibitors. The examples of inhibitor design based on the current crystal structure are given below.

Example A: Since the side chains of P<sub>3</sub> Val and P<sub>1</sub> Leu are packed against each other and there is no enzyme structure between them, cross-linking these side chains would increase the binding strength of inhibitor to memapsin 2. This is  
25 because when binding to the enzyme, the cross-linked inhibitors would have less entropy difference between the free and bound forms than their non-cross-linked counterparts (Khan, A.R., *et al.*, *Biochemistry*, 37: 16839 (1998), the teachings of which are incorporated herein in their entirety). Possible structures of the cross-linked side chains include those shown in Figure 1.

30 Example B: The same situation exists between the P<sub>4</sub> Glu and P<sub>2</sub> Asn. The current crystal structure shows that these side chains are already hydrogen bonded

-60-

to each other so the cross linking between them would also derive binding benefit as described in A. The cross-linked structures include those shown in Figure 2.

Example C: Based on the current crystal structure, the P1' Ala side chain may be extended to add new hydrophobic, Van der Waals and H-bond interactions.

5 An example of such a design is diagramed in Figure 3.

Example D: Based on the current crystal structure, the polypeptide backbone in the region of P1, P2, and P3, and the side chain of P1-Leu can be bridged into rings by the addition of two atoms (A and B in Figure 4). Also, a methyl group can be added to the beta-carbon of the P1-Leu (Figure 4).

10 Modifications and variations of the methods and materials described herein will be obvious to those skilled in the art and are intended to come within the scope of the appended claims.

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in  
15 the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

-61-

## CLAIMS

What is claimed is:

1. An inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M.
2. The inhibitor of Claim 1, comprising an isostere of the active site of memapsin 2.
3. The inhibitor of Claim 2, comprising a molecule having the general form  

$$X-L_4-P_4-L_3-P_3-L_2-P_2-L_1-P_1-L_0-P_1'-L_1'-P_2'-L_2'-P_3'-L_3'-P_4'-L_4'-Y,$$
 wherein  $P_x$  represents the substrate specificity position relative to the cleavage site which is represented by an  $-L_0-$ , and  $L_x$  represent the linking regions between each substrate specificity position,  $P_x$ , and wherein  $L_0$  is a non-hydrolyzable bond and  $P_1'$  is  $-R_1CR_3-$ , wherein  $R_1$  is a group smaller than  $CH_2OH$  (side chain of serine), and at least two other  $P$  positions are a hydrophobic group.
4. The inhibitor of Claim 1 having a  $K_i$  of less than or equal to  $10^{-6}$  M.
5. The inhibitor of Claim 1 having a  $K_i$  of less than or equal to 2 nM.
6. The inhibitor of Claim 1 having a  $K_i$  of less than or equal to 1 nM.
7. The inhibitor of Claim 1 having a root mean square difference of less than or equal to 0.5 Å for the side chain and backbone atoms for amino acids 28-441 of SEQ ID NO: 2.

-62-

8. The inhibitor of Claim 1 which is permeable to the blood brain barrier.
9. The inhibitor of Claim 1 which is less than 900 daltons in molecular weight.
10. The inhibitor of Claim 1 which blocks cleavage by memapsin 2 under  
5 physiological conditions.
11. The inhibitor of Claim 1 having a  $K_i$  of less than or equal to  $10^{-6}$  M.
12. A compound selected from the group consisting of MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-066, MMI-070, and MMI-071.
- 10 13. A compound selected from the group consisting of MMI-012, MMI-017, MMI-018, MMI-026, MMI-037, MMI-039, MMI-040, MMI-070 and MMI-071.
14. A method for treating a patient to decrease the likelihood of developing or the progression of Alzheimer's disease comprising administering to the  
15 individual an effective amount of an inhibitor of memapsin 2 selected from the group consisting of MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-070 and MMI-071.
15. The method of Claim 14, wherein the inhibitor is administered orally.
- 20 16. The method of Claim 14, wherein the inhibitor blocks cleavage of APP.
17. A method of determining the substrate side-chain preference in memapsin 2 sub-sites, comprising the steps of:
  - a) reacting a mixture of memapsin 2 substrates with memapsin 2; and

-63-

- b) determining the sub-site preference of memapsin 2 by determining relative initial hydrolysis rates of the mixture of memapsin 2 substrates.
18. The method of Claim 17, wherein the initial hydrolysis rate is determined by using a MALDI-TOF/MS device wherein the ion intensities from the MALDI-TOF/MS measurements are used for quantification of relative amounts of the substrates and the products formed thereof by hydrolysis of the substrates.
19. A method of determining the substrate side-chain preference in memapsin 2 sub-sites, comprising the steps of:
- a) preparing a combinatorial library of memapsin 2 inhibitors wherein the inhibitors comprise a base sequence taken from OM99-2;
- b) probing the library of inhibitors with memapsin 2 wherein the memapsin 2 may bind one or a plurality of inhibitors to generate one or a plurality of bound memapsin 2 ; and
- c) detecting the bound memapsin 2 with an antibody raised to memapsin 2 and an alkaline phosphatase conjugated secondary antibody.
20. The method of Claim 19, wherein the inhibitors are immobilized on beads.
21. The method of Claim 19, wherein the base sequence is EVNL\*AAEF and wherein the P<sub>3</sub>, P<sub>2</sub>, P<sub>2</sub>', and P<sub>3</sub>' sub-sites of memapsin 2 are randomized.
22. Use of an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an K<sub>i</sub> of



-64-

less than or equal to  $10^{-7}$  M for the manufacture of a medicament for the treatment of Alzheimer's disease in a human.

23. Use of an inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft, the inhibitor having an  $K_i$  of less than or equal to  $10^{-7}$  M, wherein the inhibitor has a root mean square difference of less than or equal to 0.5 Å for the side chain and backbone atoms for amino acids 18-379 of memapsin 2, for the manufacture of a medicament for the treatment of Alzheimer's disease in a human.
- 5
- 10 24. Use of a compound selected from the group consisting of MMI-005, MMI-012, MMI-017, MMI-018, MMI-025, MMI-026, MMI-037, MMI-039, MMI-040, MMI-065, MMI-066, MMI-070 and MMI-071, for the manufacture of a medicament for the treatment of Alzheimer's disease in a human.




Figure 1




Figure 2




Figure 3




Figure 4

Memapsin 2 Specificity for Positions P<sub>1</sub>'-P<sub>4</sub>'




Figure 5

gcgggagtg	gcctgccc	cgccaccag	cacggcatcc	ggctgcccc	gcgcagcggc	60
ctggggggcg	ccccctggg	gctgcggtg	ccccgggaga	ccgacgaaga	gcccagaggag	120
cccggccgga	ggggcagctt	tgtggagatg	gtggacaacc	tgaggggcaa	gtcggggcag	180
ggctactacg	tggagatgac	cgtgggcagc	ccccgcaga	cgctcaacat	cctgggtgat	240
acaggcagca	gtaactttgc	agtgggtgct	gccccccacc	ccttcctgca	tcgctactac	300
cagaggcagc	tgtccagcac	ataccgggac	ctccggaagg	gtgtgtatgt	gccctacacc	360
cagggcaagt	gggaagggga	gctgggcacc	gacctggtaa	gcatccccc	tggccccaac	420
gtcactgtgc	gtgccaacat	tgtgccatc	actgaatcag	acaagtctct	catcaacggc	480
tccaactggg	aaggcatcct	ggggctgggc	tatgctgaga	ttgccaggcc	tgacgactcc	540
ctggagcctt	tctttgactc	tctggtaaa	cagaccacg	ttcccaacct	cttctccctg	600
cagctttgtg	gtgctggctt	ccccctcaac	cagtctgaag	tgtctggcctc	tgtcggaggg	660
agcatgatca	ttggagggtat	cgaccactcg	ctgtacacag	gcagtctctg	gtatacaccc	720
atccggcggg	agtgggtatta	tgagggtgatc	attgtgcggg	tggagatcaa	tggacaggat	780
ctgaaaatgg	actgcaagga	gtacaactat	gacaagagca	ttgtggacag	tggcaccacc	840
aaccttcgtt	tgcccaagaa	agtgtttgaa	gctgcagtca	aatccatcaa	ggcagcctcc	900
tccacggaga	agtccctga	tggtttctgg	ctaggagagc	agctggtgtg	ctggcaagca	960
ggcaccaccc	cttgggaacat	tttcccagtc	atctcactct	acctaattggg	tgaggttacc	1020
aaccagtcct	tccgcatac	catacttccg	cagcaatacc	tgcggccagt	ggaagatgtg	1080
gccacgtccc	aagcagactg	ttacaagttt	gccatctcac	agtcattccac	gggcactgtt	1140
atggggagctg	ttatcatgga	gggcttctac	gttgtctttg	atcgggcccc	aaaacgaatt	1200
ggctttgtctg	tcagcgcttg	ccatgtgcac	gatgagttca	ggacggcagc	ggtggaaggc	1260
ccttttgtca	ccttggacat	ggaagactgt	ggctacaaca	ttccacagac	agatgagtca	1320
accctcatga	ccatagccta	tgtcatggct	gccatctgcg	ccctcttcat	gctgccactc	1380
tgcctcatgg	tgtgtcagtg	gcgctgcctc	cgctgcctgc	gccagcagca	tgatgacttt	1440
gctgatgaca	tctccctgct	gaagtggaga	ggcccatggg	cagaagatag	agattcccc	1500
ggaccacacc	tccgtggttc	actttggtca	caagtaggag	acacagatgg	cacctgtggc	1560
cagagcacct	caggaccctc	cccaccaccc	aatgcctct	gccttgatgg	agaaggaaaa	1620
ggctggcaag	gtgggttcca	gggactgtac	ctgtaggaaa	cagaaaagag	aagaaaagag	1680
cactctgtctg	gcgggaatac	tcttggtcac	ctcaaattta	agtcgggaaa	ttctgtgct	1740
tgaaacttca	gcccgaacc	tttgtccacc	attcctttta	attctccaac	ccaaagtatt	1800
cttcttttct	tagtttcaga	agtactggca	tcacacgcag	gttaccttgg	cggtgtgtccc	1860
tgtggtagcc	tggcagagaa	gagaccaagc	ttgtttccct	gctggccaaa	gtcagtagga	1920
gaggatgcac	agtttgctat	ttgctttaga	gacagggact	gtataaaca	gcctaaccatt	1980
ggtgcaagca	ttgcctcttg	aattaaaaaa	aaactagatt	gactatttat	acaaatgggg	2040
ggcgctggaa	agaggagaag	gagagggagt	acaaagacag	ggaatagtgg	gatcaaagct	2100
aggaaaggca	gaaacacaac	cactcaccag	tcctagtttt	agacctcatc	tccaagatag	2160
catcccatct	cagaagatgg	gtgttggttt	caatgttttc	ttttctgtgg	ttgcagcctg	2220
accaaagtg	agatgggaag	ggcttatcta	gccaaagagc	tcttttttag	ctctcttaaa	2280
tgaagtggcc	actaagaagt	tocacttaac	acatgaattt	ctgccatatt	aatttcattg	2340
tctctatctg	aaccaccctt	tattctacat	atgataggca	gcactgaaat	atcctaacc	2400
cctaagctcc	aggtgcccctg	tgggagagca	actggactat	agcagggctg	ggctctgtct	2460
tcttggtcat	aggtcactc	tttcccccaa	atcttctctt	ggagctttgc	agccaagggtg	2520
ctaaaaggaa	taggtaggag	acctcttcta	tctaattcctt	aaaagcataa	tgttgaacat	2580
tcattcaaca	gctgatgcc	tataaccctt	gcctggattt	cttctatta	ggctataaga	2640
agtagcaaga	tctttacata	attcagagtg	gtttcattgc	cttctaccc	tctctaattg	2700
cccctccatt	tatttgacta	aagcatcrca	cagtggcact	agcattatac	caagagtatg	2760
agaaatacag	tgttttatgg	ctctaaccatt	actgccttca	gtatcaaggc	tgccctggaga	2820
aaggatggca	gcctcagggc	ttccttatgt	cctccaccac	aagagctcct	tgatgaagg	2880
catctttttc	ccctatcctg	ttcttccctt	ccccgctcct	aatggtagct	gggtaccag	2940
gctggttctt	gggctaggta	gtggggacca	agttcattac	ctccctatca	gttctagcat	3000
agtaaaactac	ggtaccagtg	ttagtgggaa	gagctgggtt	ttcctagtat	acccactgca	3060
tcctactcct	acctggtcaa	ccgctgctt	ccaggatagg	gacctgctaa	gtgtggaatt	3120
acctgataag	ggagagggaa	atacaaggag	ggcctctgg	gttcttgcc	tcagccagct	3180
gcccmaagc	cataaaccaa	taaamcaaga	atactgagtc	taaaaaaaaa	aaaaaaaaaaa	3240
aaaaaaaaaa	aa					3252

Figure 6

Ala	Gly	Val	Leu	Pro	Ala	His	Gly	Thr	Gln	His	Gly	Ile	Arg	Leu	Pro	1	5	10	15
Leu	Arg	Ser	Gly	Leu	Gly	Gly	Ala	Pro	Leu	Gly	Leu	Arg	Leu	Pro	Arg	20	25	30	
Glu	Thr	Asp	Glu	Glu	Pro	Glu	Glu	Pro	Gly	Arg	Arg	Gly	Ser	Phe	Val	35	40	45	
Glu	Met	Val	Asp	Asn	Leu	Arg	Gly	Lys	Ser	Gly	Gln	Gly	Tyr	Tyr	Val	50	55	60	
Glu	Met	Thr	Val	Gly	Ser	Pro	Pro	Gln	Thr	Leu	Asn	Ile	Leu	Val	Asp	65	70	75	80
Thr	Gly	Ser	Ser	Asn	Phe	Ala	Val	Gly	Ala	Ala	Pro	His	Pro	Phe	Leu	85	90	95	
His	Arg	Tyr	Tyr	Gln	Arg	Gln	Leu	Ser	Ser	Thr	Tyr	Arg	Asp	Leu	Arg	100	105	110	
Lys	Gly	Val	Tyr	Val	Pro	Tyr	Thr	Gln	Gly	Lys	Trp	Glu	Gly	Glu	Leu	115	120	125	
Gly	Thr	Asp	Leu	Val	Ser	Ile	Pro	His	Gly	Pro	Asn	Val	Thr	Val	Arg	130	135	140	
Ala	Asn	Ile	Ala	Ala	Ile	Thr	Glu	Ser	Asp	Lys	Phe	Phe	Ile	Asn	Gly	145	150	155	160
Ser	Asn	Trp	Glu	Gly	Ile	Leu	Gly	Leu	Ala	Tyr	Ala	Glu	Ile	Ala	Arg	165	170	175	
Pro	Asp	Asp	Ser	Leu	Glu	Pro	Phe	Phe	Asp	Ser	Leu	Val	Lys	Gln	Thr	180	185	190	
His	Val	Pro	Asn	Leu	Phe	Ser	Leu	Gln	Leu	Cys	Gly	Ala	Gly	Phe	Pro	195	200	205	
Leu	Asn	Gln	Ser	Glu	Val	Leu	Ala	Ser	Val	Gly	Gly	Ser	Met	Ile	Ile	210	215	220	
Gly	Gly	Ile	Asp	His	Ser	Leu	Tyr	Thr	Gly	Ser	Leu	Trp	Tyr	Thr	Pro	225	230	235	240

Figure 7A

Ile	Arg	Arg	Glu	Trp	Tyr	Tyr	Glu	Val	Ile	Ile	Val	Arg	Val	Glu	Ile	245	250	255
Asn	Gly	Gln	Asp	Leu	Lys	Met	Asp	Cys	Lys	Glu	Tyr	Asn	Tyr	Asp	Lys	260	265	270
Ser	Ile	Val	Asp	Ser	Gly	Thr	Thr	Asn	Leu	Arg	Leu	Pro	Lys	Lys	Val	275	280	285
Phe	Glu	Ala	Ala	Val	Lys	Ser	Ile	Lys	Ala	Ala	Ser	Ser	Thr	Glu	Lys	290	295	300
Phe	Pro	Asp	Gly	Phe	Trp	Leu	Gly	Glu	Gln	Leu	Val	Cys	Trp	Gln	Ala	305	310	315
Gly	Thr	Thr	Pro	Trp	Asn	Ile	Phe	Pro	Val	Ile	Ser	Leu	Tyr	Leu	Met	325	330	335
Gly	Glu	Val	Thr	Asn	Gln	Ser	Phe	Arg	Ile	Thr	Ile	Leu	Pro	Gln	Gln	340	345	350
Tyr	Leu	Arg	Pro	Val	Glu	Asp	Val	Ala	Thr	Ser	Gln	Asp	Asp	Cys	Tyr	355	360	365
Lys	Phe	Ala	Ile	Ser	Gln	Ser	Ser	Thr	Gly	Thr	Val	Met	Gly	Ala	Val	370	375	380
Ile	Met	Glu	Gly	Phe	Tyr	Val	Val	Phe	Asp	Arg	Ala	Arg	Lys	Arg	Ile	385	390	395
Gly	Phe	Ala	Val	Ser	Ala	Cys	His	Val	His	Asp	Glu	Phe	Arg	Thr	Ala	405	410	415
Ala	Val	Glu	Gly	Pro	Phe	Val	Thr	Leu	Asp	Met	Glu	Asp	Cys	Gly	Tyr	420	425	430
Asn	Ile	Pro	Gln	Thr	Asp	Glu	Ser	Thr	Leu	Met	Thr	Ile	Ala	Tyr	Val	435	440	445
Met	Ala	Ala	Ile	Cys	Ala	Leu	Phe	Met	Leu	Pro	Leu	Cys	Leu	Met	Val	450	455	460
Cys	Gln	Trp	Arg	Cys	Leu	Arg	Cys	Leu	Arg	Gln	Gln	His	Asp	Asp	Phe	465	470	475
Ala	Asp	Asp	Ile	Ser	Leu	Leu	Lys									485		

Figure 7B

Met	Ala	Ser	Met	Thr	Gly	Gly	Gln	Gln	Met	Gly	Arg	Gly	Ser	Met	Ala
1				5					10					15	
Gly	Val	Leu	Pro	Ala	His	Gly	Thr	Gln	His	Gly	Ile	Arg	Leu	Pro	Leu
			20					25					30		
Arg	Ser	Gly	Leu	Gly	Gly	Ala	Pro	Leu	Gly	Leu	Arg	Leu	Pro	Arg	Glu
		35					40					45			
Thr	Asp	Glu	Glu	Pro	Glu	Glu	Pro	Gly	Arg	Arg	Gly	Ser	Phe	Val	Glu
	50					55					60				
Met	Val	Asp	Asn	Leu	Arg	Gly	Lys	Ser	Gly	Gln	Gly	Tyr	Tyr	Val	Glu
65					70					75					80
Met	Thr	Val	Gly	Ser	Pro	Pro	Gln	Thr	Leu	Asn	Ile	Leu	Val	Asp	Thr
				85					90					95	
Gly	Ser	Ser	Asn	Phe	Ala	Val	Gly	Ala	Ala	Pro	His	Pro	Phe	Leu	His
			100					105					110		
Arg	Tyr	Tyr	Gln	Arg	Gln	Leu	Ser	Ser	Thr	Tyr	Arg	Asp	Leu	Arg	Lys
		115					120					125			
Gly	Val	Tyr	Val	Pro	Tyr	Thr	Gln	Gly	Lys	Trp	Glu	Gly	Glu	Leu	Gly
	130					135					140				
Thr	Asp	Leu	Val	Ser	Ile	Pro	His	Gly	Pro	Asn	Val	Thr	Val	Arg	Ala
145					150					155					160
Asn	Ile	Ala	Ala	Ile	Thr	Glu	Ser	Asp	Lys	Phe	Phe	Ile	Asn	Gly	Ser
				165					170					175	
Asn	Trp	Glu	Gly	Ile	Leu	Gly	Leu	Ala	Tyr	Ala	Glu	Ile	Ala	Arg	Pro
		180						185					190		
Asp	Asp	Ser	Leu	Glu	Pro	Phe	Phe	Asp	Ser	Leu	Val	Lys	Gln	Thr	His
		195					200					205			
Val	Pro	Asn	Leu	Phe	Ser	Leu	Gln	Leu	Cys	Gly	Ala	Gly	Phe	Pro	Leu
	210					215					220				
Asn	Gln	Ser	Glu	Val	Leu	Ala	Ser	Val	Gly	Gly	Ser	Met	Ile	Ile	Gly
225					230					235					240

Figure 8A



Gly	Ile	Asp	His	Ser	Leu	Tyr	Thr	Gly	Ser	Leu	Trp	Tyr	Thr	Pro	Ile	245	250	255
Arg	Arg	Glu	Trp	Tyr	Tyr	Glu	Val	Ile	Ile	Val	Arg	Val	Glu	Ile	Asn	260	265	270
Gly	Gln	Asp	Leu	Lys	Met	Asp	Cys	Lys	Glu	Tyr	Asn	Tyr	Asp	Lys	Ser	275	280	285
Ile	Val	Asp	Ser	Gly	Thr	Thr	Asn	Leu	Arg	Leu	Pro	Lys	Lys	Val	Phe	290	295	300
Glu	Ala	Ala	Val	Lys	Ser	Ile	Lys	Ala	Ala	Ser	Ser	Thr	Glu	Lys	Phe	305	310	315
Pro	Asp	Gly	Phe	Trp	Leu	Gly	Glu	Gln	Leu	Val	Cys	Trp	Gln	Ala	Gly	325	330	335
Thr	Thr	Pro	Trp	Asn	Ile	Phe	Pro	Val	Ile	Ser	Leu	Tyr	Leu	Met	Gly	340	345	350
Glu	Val	Thr	Asn	Gln	Ser	Phe	Arg	Ile	Thr	Ile	Leu	Pro	Gln	Gln	Tyr	355	360	365
Leu	Arg	Pro	Val	Glu	Asp	Val	Ala	Thr	Ser	Gln	Asp	Asp	Cys	Tyr	Lys	370	375	380
Phe	Ala	Ile	Ser	Gln	Ser	Ser	Thr	Gly	Thr	Val	Met	Gly	Ala	Val	Ile	385	390	395
Met	Glu	Gly	Phe	Tyr	Val	Val	Phe	Asp	Arg	Ala	Arg	Lys	Arg	Ile	Gly	405	410	415
Phe	Ala	Val	Ser	Ala	Cys	His	Val	His	Asp	Glu	Phe	Arg	Thr	Ala	Ala	420	425	430
Val	Glu	Gly	Pro	Phe	Val	Thr	Leu	Asp	Met	Glu	Asp	Cys	Gly	Tyr	Asn	435	440	445
Ile	Pro	Gln	Thr	Asp	Glu	Ser	Thr	Leu	Met	Thr	Ile	Ala	Tyr	Val	Met	450	455	460
Ala	Ala	Ile	Cys	Ala	Leu	Phe	Met	Leu	Pro	Leu	Cys	Leu	Met	Val	Cys	465	470	475
Gln	Trp	Arg	Cys	Leu	Arg	Cys	Leu	Arg	Gln	Gln	His	Asp	Asp	Phe	Ala	485	490	495
Asp	Asp	Ile	Ser	Leu	Leu	Lys										500		

Figure 8B

1 MAQALPWLLL WMGAGVLP AH GTQHGIRLPL RSGLGGA PLG LRLPRETDEE  
51 PEEPGRRG SF VEMVDNLRGK SGQGYVEMT VGSP PQTLNI LVDTGSSNFA  
101 VGAAPHPFLH RYYQRQLSST YRDLRKG VYV PYTQ GKWEGE LGTDLVSI PH  
151 GPNVTVRANI AAITESDKFF INGSNWEGIL GLAYAEIARP DDSLEPFFDS  
201 LVKQTHVPNL FSLQLCGAGF PLNQSEVLAS VGGSMIIGGI DHSLYTGSLW  
251 YTPIRREWYY EVIIVRVEIN GQDLKMDCKE YNYDKSIVDS GTTNLRLPKK  
301 VFEEAVKSIK AASSTEKFPD GFWLGEQLVC WQAGTTPWNI FPVISLYLMG  
351 EVTNQSFRIT ILPQQYLRPV EDVATSQDDC YKFAISQSST GTVMGAVIME  
401 GFYVVFDRAR KRIGFAVSAC HVHDEFRTAA VEGPFVTLDM EDCGYNIPQT  
451 DESTLMTIAY VMAAICALFM LPLCLMVCQW RCLRCLRQQH DDFADDISLL  
501 K

Figure 9

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**